

NEW neat STUFF

mtDNA KIT

- Efficient isolation of mtDNA from whole blood, cell culture, or tissue
- Results in ~ 90 minutes
- No Hazardous Reagents

CYTOTOXICITY

- Simple 2 step procedure
- Results in ~ 5 minutes
- No Pre-Treatment, No Dye Preparation, and No Hazardous Waste

MICROENCAPSULATED OsO_4 & $Sc(OTf)_3$

- Popular Catalysts with Improved Safety
- Low Toxicity
- Easy Recovery and Recyclable

HPLC WAKOSIL II PACKING

- Symmetrical spherical packing for HPLC, flash & column chromatography
- High Purity and High Efficiency
- High Lot to Lot Reproducibility

SLP REAGENT SET

- Highly sensitive peptidoglycan and (1-3)-B-glucan Assay.
- Detection of Microorganism Contamination
- No Instrument Required

TAPS-SULFONATE

- Novel reagent for S-Alkylsulfidation
- Solubilizes denatured proteins and allows purification prior to refolding
- High Efficiency and High Recovery

From your favorite supplier of
Anti-Asialo, Lys-C, Digitonin,
DNA Kits, and Trichostatin A

Wako

Wako USA Wako GmbH
P: (877) 714-1920 49 (02131) 3111-0
F: (804) 271-7791 49 (02131) 311100
www.wako-chem.co.jp

Subsidiaries of
Wako Pure Chemical Industries, Ltd. Osaka Japan

SCIENCE'S COMPASS

change in preformed nutrients [resulting in a cumulative decrease in deep-water phosphate of about 0.02 micromoles per kilogram ($\mu\text{mol/kg}$) over 20 years and a cumulative increase in deep-water nitrate of 0.14 $\mu\text{mol/kg}$ over the same period (2)] could also explain the reported N:P trends. Such a change in preformed nutrients, caused for example by a slight shift in the deep-water source regions rather than in biology, would be undetectable with the presently available data because of large seasonal variability and inadequate sampling.

For North Pacific deep waters, Pahlow and Riebesell report a small increase in AOU and suggest enhanced export production as a cause. However, increasing AOU could also be caused by decreasing water ventilation rates—a change predicted as a response to global warming (3) and perhaps manifested in recent chlorofluorocarbon budgets (4). Testing this alternative ventilation mechanism poses a formidable (and so far open) challenge.

There are many observations that demonstrate that the marine biota is variable, especially on local scales (5). However, the conclusions by Pahlow and Riebesell that the marine biota has changed over large spatial and temporal scales are unsubstantiated because alternative hypotheses cannot be firmly excluded.

Nicolas Gruber

Institute of Geophysics and Planetary Physics and Department of Atmospheric Sciences, University of California, Los Angeles, Los Angeles, CA 90095, USA

Klaus Keller

Robert M. Key

Department of Geosciences, Princeton University, Guyot Hall, Princeton, NJ 08544, USA

References and Notes

1. L. I. Gordon *et al.*, *Eos Trans. Am. Geophys. Union* **80** (suppl.), OS45 (1999); C. W. Mordy *et al.*, *Eos Trans. Am. Geophys. Union* **80** (suppl.), OS43 (1999).
2. Computed by using the rates of changes reported by Pahlow and Riebesell and their equation 3, solved for the change in nutrient concentration ($n_t - n_0$) and adopting a value of 120 $\mu\text{mol/kg}$ for AOU₀.
3. S. Manabe *et al.*, *J. Climate* **4**, 785 (1991); J. L. Sarmiento *et al.*, *Nature* **393**, 245 (1998).
4. W. S. Broecker *et al.*, *Science* **286**, 132 (1999).
5. D. M. Karl *et al.*, *Nature* **373**, 230 (1995).

Response

Gruber, Keller, and Key suggest alternative explanations for the trends in deep-ocean Redfield ratios obtained in our analysis. They suggest that a small decrease in preformed nutrients could explain the trends observed in the North Atlantic and would be undetectable with presently available data. As we outlined in our report, however, the decrease in preformed phosphate concentrations necessary to generate the observed increase in the N:P ratio would have to be at least 12% or about 0.2 micromolar over 20 years. A change in preformed phosphate concentrations of this magni-

tude, should it exist, could be easily detected in the data available for the time period covered in our analysis.

Gruber *et al.* say that increased aeolian nitrogen (N) deposition "would decrease the ratio of apparent oxygen utilization (AOU) to nitrate" in the North Atlantic. This would be true only if primary production in the North Atlantic was not limited by nitrogen. Although a transition from nitrogen to phosphorus limitation has been suggested for some areas, nitrogen is considered to be the dominant limiting nutrient in the North Atlantic (1). A recent study by Wu *et al.* (2) reports phosphate depletion in parts of the western North Atlantic. The authors conclude, however, that the concentration of available nitrogen is decreased to limiting levels in these regions because of the formation of refractory dissolved organic nitrogen. Although increasing aeolian nitrogen deposition may have in fact reduced the AOU:N ratio locally in phosphorus-limited regions, the effect would not be detected by our large-scale analysis.

For the North Pacific, Gruber *et al.* suggest that decreasing deep-water ventilation rates as a result of global warming may explain the observed increase in AOU. North Pacific deep waters originate almost entirely from the South Pacific (3). If decreasing deep-water ventilation had caused the trends indicated for the North Pacific, similar trends should also have occurred in the South Pacific. Because this is not observed, changes in deep-water ventilation are not likely to be the main cause for the changes in AOU and oxidative ratios observed in the North Pacific.

In essence, although the nonbiologically mediated processes suggested by Gruber *et al.* may have contributed to the observed temporal trends in deep-ocean Redfield ratios, they cannot account for their magnitude. The changes in marine biota that we proposed in our report thus remain the most plausible explanation for the observed trends.

Markus Pahlow*

Ulf Riebesell

Alfred Wegener Institute for Polar and Marine Research, Post Office Box 120161, D-27515 Bremerhaven, Germany

*Present address: Bedford Institute of Oceanography, Dartmouth, Nova Scotia, B2Y 4A2, Canada

References

1. K. A. Fanning, *Nature* **339**, 469 (1989).
2. J. Wu, W. Sunda, E. A. Boyle, D. M. Karl, *Science* **289**, 759 (2000).
3. A. M. Macdonald, *Prog. Oceanogr.* **41**, 281 (1998).

CORRECTIONS AND CLARIFICATIONS

News Focus: "Louisiana's vanishing wetlands: going, going,..." by J. Bourne (15 Sept., p.1860). The word "levy" and its derivatives were used in the article instead of "levee."