

- phys. 349, 376 (1998).
 2. M. Levine et al., *Proc. Natl. Acad. Sci. U.S.A.* 93, 3704 (1996).
 3. A. Snyder, L. Skoza, Y. Kikkawa, *Lung* 161, 111 (1983).

IN RESPONSE TO BAR-OR, WE ACCEPT

the criticism that this study was conducted at pH 7.0 rather than pH 7.4. We were mainly concerned with eliminating transition metal ions and with conducting the reactions at neutral pH to show that we were not simply observing the well-known transition metal ion pro-oxidant effect (1). Clearly, the only way to prove that the lipid hydroperoxide-derived bifunctional electrophiles are formed at intracellular pH is to conduct both in vitro cell culture and in vivo experiments. From a theoretical perspective, there is no reason to suppose that a pH difference of 0.4 units would affect the reaction that we discovered. Thus, at pH 7.0 and pH 7.4, both vitamin C and polyunsaturated fatty acid hydroperoxides exist almost exclusively in their corresponding anion forms. Furthermore, small changes in pH have essentially no effect on the reactions of lipid hydroperoxide-derived bi-

functional electrophiles with DNA. Below pH 6.0, 4,5-epoxy-2(E)decenal is unstable. However, this is well below the pH that we used in our reactions. Subsequent to the submission of our report, we tested the reaction of vitamin C with lipid hydroperox-

"From a theoretical perspective, there is no reason to suppose that a pH difference of 0.4 units would affect the reaction that we discovered."

ides at pH 7.4 and found that the product profile was essentially the same as that observed at pH 7.0 (2).

It is certainly conceivable that DNA damage caused by acidosis results from Cu/ascorbate-mediated radical formation rather than ascorbate alone. However, our study showed that lipid radicals could also be formed by the Cu/ascorbate system. This

would result in generation of the same lipid hydroperoxide-derived bifunctional electrophiles that we reported. Furthermore, the mutagenic potential of the resulting DNA lesions, such as etheno-2'-deoxyadenosine (3), is much greater than that resulting from oxidative damage (3, 4). This suggests that it will be important in the future to monitor human populations for DNA lesions that result from lipid hydroperoxide-derived genotoxins, as we suggested in our report.

SEON HWA LEE, TOMOYUKI OE, IAN A. BLAIR*

Center for Cancer Pharmacology, 1254 BRB II/III, 421 Curie Boulevard, University of Pennsylvania, Philadelphia, PA 19104-6160, USA

*To whom correspondence should be addressed.

E-mail: ian@spirit.gcrc.upenn.edu

References and Notes


1. A. R. Prottogente, A. Rehman, B. Halliwell, C. A. Rice-Evans, *Biochem. Biophys. Res. Commun.* 277, 535 (2000).
2. S. H. Lee, T. Oe, I. A. Blair, unpublished observations.
3. R. L. Levine et al., *Cancer Res.* 60, 4098 (2000).
4. J. Cadet et al., *Mutat. Res.* 424, 9 (1999).

Sustainability: Insight from Industrial Ecology

SUSTAINABILITY IS ONE OF THE MOST important issues we face today, and thus it

Discounted Scientific and Technical Titles for Members at the AAAS Bookstore

Update your library at:
www1.fatbrain.com/AAAS



visit our website and register to

WIN

a New
Volkswagen
Beetle from...

spectrum
Laboratory Products
An ISO 9002 Registered Company

1.800.632.9154



ScientistsWanted.com

Circle No. 55 on Readers' Service Card

is essential that it be addressed using the most intelligent and creative methods available. In their Policy Forum "Sustainability science" (Science's Compass, 27 Apr., p. 641), Robert W. Kates and co-authors seem to suggest applying natural systems methodology—that is, an observational approach—to human systems. Indeed, studying nature allows us to capture the problem of sustainability, but it gives us no insight into the solution. We suggest complementing the observational approach with a more active, solution-driven approach.

A conceptualization of the solution-driven approach is embodied in the field of industrial ecology (IE), which has been defined as the "science of sustainability" (1). The systems approaches of IE help in understanding and minimizing the burdens imposed by society on nature through methods such as material and energy flow studies, dematerialization and decarbonization, life-cycle assessment, design for the environment, product stewardship, eco-industrial parks, Earth systems engineering/management, product-oriented environmental policy, and eco-efficiency. Some of the approaches are technical; some involve policy. In concert with sustainability science, IE also aims, as Kates *et al.* say, to "understand the fundamental character of interactions between nature and society."

Since its beginnings in the late 1980s, the IE community has been interdisciplinary, including engineers, physicists,

"The systems approaches of [industrial ecology] help in understanding and minimizing the burdens imposed by society on nature..."

ecologists, economists, sociologists, and other scientists working in government, academia, and industry. Two Gordon Conferences, *The Journal of Industrial Ecology*, and the recent formation of the International Society for Industrial Ecology have played a major role in crystallizing the "science of sustainability." We can all benefit from the synergism of many minds. In the spirit of cooperation, we invite the nascent sustainability science community into a dialog with our community. We see the two groups as proposing complementary and, in some cases, simi-

lar pathways toward a scientific approach to sustainability.

BARBARA PARISH KARN,* DIANA BAUER†

U.S. Environmental Protection Agency, Office of Research and Development, National Center for Environmental Research, Washington, DC 20460, USA

*To whom correspondence should be addressed. E-mail: karn.barbara@epa.gov

†AAAS fellow

References and Notes

1. T. Graedel, *Environ. Sci. Technol.* **34**, 28A (2000).

Separate Identity, Separate Career

WE WERE HONORED BY THE NEWS FOCUS article on our careers in physics and astrophysics ("Twin stars of astrophysics make room for two" by M. Sincell, 10 Aug., p. 1040). The article itself illustrates some of the challenges we have faced as identical twins attempting to forge separate professional identities and overcome the distraction that our special relationship creates for others.

Not only were our identities reversed in the caption of the photograph that accompanied the article, but space was devoted to describing several twin incidents at the cost of barely mentioning the individual scientific contributions of one of us (FKL). We would like to mention a few highlights for the record.

In early 1972, FKL proposed that the recently discovered x-ray pulsars are accreting magnetic neutron stars. With his Illinois colleagues and students, he developed the theory of such stars. With his student Pranab Ghosh, he also developed the standard theory of disk accretion by spinning magnetic stars. FKL not only played an important role in the conception, development, and operation of NASA's Rossi X-Ray Timing Explorer mission, but also, with his students, developed a theory of the kilohertz brightness oscillations of neutron stars and black holes that were discovered with the Rossi Explorer. He and his students have shown that these oscillations can be used to determine how matter and radiation move in the strong gravitational fields near neutron stars and black holes and to constrain the properties of the ultradense matter in neutron stars.

FREDERICK K. LAMB,¹* DON Q. LAMB²

¹University of Illinois at Urbana-Champaign, Departments of Physics and Astronomy and Center for Theoretical Astrophysics, Urbana, IL 61801, USA. ²University of Chicago, Department of Astronomy and Astrophysics, Chicago, IL 60637, USA

*To whom correspondence should be addressed.

E-mail: f-lamb@uiuc.edu

The State of Coral Reef Science

IN THEIR REPORT "REGIONAL-SCALE ASSEMBLY rules and biodiversity of coral reefs" (25 May, p. 1532), David R. Bellwood and Terry P. Hughes make several statements that warrant rebuttal. First, they say that "our understanding of regional-scale patterns of biodiversity on coral reefs...has changed little since the seminal work of Stehli and Wells..." This comment disregards a 30-year-long research program conducted by the Australian Institute of Marine Science (AIMS), which effectively follows on from where Stehli and Wells (1971) left off (1). Second, the authors say that "the processes that shape [biodiversity gradient] patterns [on coral reefs] remain elusive," yet the geological background and the role of ocean currents in dispersion, genetic connectivity, temperature, and habitat variation have all been well addressed in more than 100 publications. And third, Bellwood and Hughes suggest that their work is the "first" to quantify the "biogeographic variation in species composition...along biodiversity gradients." I spent 20 years doing precisely that for corals.

Since the late 19th century, fish and coral data have led the marine contribution to biogeographic and evolutionary debate. Now, at least 90% of species likely to be encountered on any reef have been described and their distribution ranges recorded. This information is available in Geographic Information Systems (GIS) databases, which permit the sorts of sophisticated analyses that support terrestrial biogeography. Of the 31 references in Bellwood and Hughes' report, 23 were published in the 1980s or earlier, and all (including those of the present author) have been superseded, both taxonomically and biogeographically. The major marine conservation organizations, working with GIS databases (of reefs, corals, fish, and molluscs), are going to considerable lengths to preserve both credibility and accuracy (2), but Bellwood and Hughes appear to have drawn little upon these extensive resources.

J. E. N. VERON

Chief Scientist, Australian Institute of Marine Science, PMB 3, Townsville, MC 4810, Australia. E-mail: jveron@aims.gov.au

References and Notes

1. The wealth of coral biogeographic publications that have changed every aspect of this thesis is summarized in J. E. N. Veron, *Corals in Space and Time: the Biogeography and Evolution of the Scleractinia* (Comstock, Ithaca, NY, 1995), which received the Australian National Marine Science Award for 1996.
2. For example, high-resolution biogeographic studies that support marine conservation will be highlighted in a forthcoming World Heritage Tropical Marine Biodiversity Assessment Workshop sponsored by the UN, UNESCO, NOAA, the World Conservation Monitoring Centre, and IUCN.