

tradict his argument when noting that such relicts have accumulated "over vast periods of time.") Many of our centers of endemism appear to be areas of evolutionary innovation (1). Other things being equal, restricted-range species are more vulnerable to extinction than widespread species and require focused conservation effort.

Our measure of range size (number of grid cells with reef habitat in a species' range) could lead to larger centers of endemism being defined in regions where reefs are sparse, compared with regions where they are more extensive, but our results show that this did not happen. We give equal weight to all four taxa and do not dismiss corals, as suggested by Baird *et al.* Not all centers of endemism are rich in endemics for all taxa, but conservation efforts in them must protect reefs as ecosystems, including all species present.

Threats to reefs constitute a key part of our analysis. The Great Barrier Reef lies within the world's largest marine park and so does not qualify as a hotspot requiring urgent conservation effort. We presume that it is already receiving the attention it needs. We emphasize the importance of integrating terrestrial and marine conservation, because many of the threats facing reefs originate inland. Although protection of Lord Howe's

reefs will clearly not depend on land use in distant New Zealand, conservation of the Great Barrier Reef will not succeed without addressing land-based threats. Nor would it be effective in many of the other regions where terrestrial and marine hotspots abut, such as the Philippines or Southern Mascarene Islands. Finally, efforts to mitigate climate change are needed to secure the future of all the world's reefs, but place-based initiatives must get under way even as the world warms. Our findings point to the most pressing priorities and, in spreading conservation action among many places, should ultimately leave us with more options (and species) in a warmer world than if conservation were concentrated only in the richest regions.

CALLUM M. ROBERTS,<sup>1</sup> COLIN J. MCCLEAN,<sup>1</sup>

JOHN E. N. VERON,<sup>2</sup> JULIE P. HAWKINS,<sup>1</sup>

GERALD R. ALLEN,<sup>3,7</sup> DON E. MCALLISTER,<sup>4\*</sup>

CRISTINA G. MITTERMEIER,<sup>3</sup> FREDERICK W. SCHUELER,<sup>5</sup>

MARK SPALDING,<sup>6</sup> FRED WELLS,<sup>7</sup> CARLY VYNNE,<sup>3</sup>

TIMOTHY B. WERNER<sup>3</sup>

<sup>1</sup>Environment Department, University of York, York, YO10 5DD, UK. <sup>2</sup>Australian Institute of Marine Science, Private Mail Box 3, Townsville Mail Center, Queensland 4810, Australia. <sup>3</sup>Center for Applied Biodiversity Science, Conservation International, 1919 M Street NW, Suite 600, Washington, DC 20036, USA. <sup>4</sup>Ocean Voice International, Post Of-

fice Box 20060, Regional Post Office Perth Mews, Perth, Ontario, Canada K7H 3M6. <sup>5</sup>Eastern Ontario Biodiversity Museum, Box 1860, Kemptville, Ontario, Canada K0G 1J0. <sup>6</sup>United Nations Environment Program—World Conservation Monitoring Centre, 219 Huntingdon Road, Cambridge, CB3 0DL, UK. <sup>7</sup>Western Australian Museum, Francis Street, Perth, Western Australia 6000, Australia.

\*Deceased.

#### References and Notes

1. C. R. C. Sheppard, A. R. G. Price, C. M. Roberts, *Marine Ecology of the Arabian Region* (Academic Press, London, 1992).

#### CORRECTIONS AND CLARIFICATIONS



**RANDOM SAMPLES:** "Plastic invasion" (26 April 2002, p. 653): The study described analyzed roughly 6000 pieces of beach debris, not 200, and Antarctic waters (not the continent) are predicted to warm 2°C over the next century.

**LETTERS:** "Another look at MgB<sub>2</sub> and YBCO wires" by R. A. Hawsey and D. E. Peterson (26 April, p. 655). The penalty in refrigeration resulting from a reduction in operating temperature from 68 to 25 K is a factor of 3 to 4, not 3 to 4% as stated in the letter.

## MILLIPORE

# pure BACceleration

Map genomes faster by cutting BAC prep time in half with the Montage™ BAC96 Miniprep kit. Finally, a 96-well purification kit optimized for BACs. Prepare high quality BAC DNA in just 60 minutes. No isopropanol. No centrifugation. You'll also get longer read lengths, so your mapping goes even faster. To qualify for a sample kit, visit [www.millipore.com/montageBAC](http://www.millipore.com/montageBAC).



In the US and Canada, call  
1-800-MILLIPORE.  
In Europe, fax +33-3.88.38.91.95.  
In Japan, call (03)5442-9719.

© 2002 Millipore Corporation. All rights reserved. Millipore is a registered trademark and Montage is a trademark of Millipore Corporation.

Stop by booths 2814 and 2815 at Pittcon