

No nukes? Sonoluminescent bubbles look bad for fusion.

'We understand what's going on inside the bubble,' and if this is what you believe the science is, you should be suspicious of the Taleyarkhan paper," says Crum. The authors of that paper could not be reached for comment. But for now, at least, skeptical voices rule the acoustic waves.

initiate fusion.

"They're saying,


-CHARLES SEIFE

CLIMATE PREDICTION

Signs of Success in Forecasting El Niño

El Niño, the sleeping giant of climate, awakened earlier this month, according to government scientists. Six months ago, those same researchers went out on a limb when they recognized stirrings around the Pacific as likely signs that El Niño's warming of tropical waters would soon return after a 4-year absence. Their early-January forecast appears to be holding up; if the warming trend continues, El Niño's often disastrous weather shifts around the globe should crest around the end of the year. A full year's warning of El Niño's peak would be much better than forecasters achieved last time. But soon will come the next test: Will this El Niño develop into a weak-to-moderate warming this winter, as now predicted, or another barnburner like last time?

El Niño forecasting has a long and



Good call. Early this year, government scientists correctly predicted an El Niño warming while the tropical Pacific was still near normal.

checkered history. In the 1960s and even early '70s, monitoring of the tropical Pacific was so spotty that full-blown El Niños could pop up around Christmastime without warning. A simple model of tropical Pacific winds and currents successfully predicted a

www.cpc.ncep.noaa.gov/products/analysis_ monitoring/enso_advisory/index.html

† iri.columbia.edu/climate/ENSO/currentinfo/ OuickLook.html

iri.columbia.edu/climate/ENSO/currentinfo/ SST_table.html

Pacific warming for the first time in 1986, but forecasters' optimism was short-lived. It was only with the 1997 super-El Niño that human and computer forecasters had some measure of success, and even then they were criticized for a months-late alert and gross underestimation of the event's huge scale (Science, 13 October 2000, p. 257).

This time around, some forecasters seem to have gotten the onset of El Niño right. At the National Weather Service's Climate Prediction Center (CPC) in Camp Springs, Maryland, meteorologist Vernon Kousky and a half-dozen colleagues put out a monthly "diagnostic discussion." Their Web site report* sorts through observations from ships, islands, satellites, and buoys across the Pacific and evaluates forecasts from more than a dozen models run by CPC and others. Last fall, while most of the tropical Pacific was at near-normal temperatures, the CPC group started talking about a warming trend that would likely continue into 2002, although as a group the models dithered from month to month between calling for normal and somewhat warmer conditions.

On 9 January, while the crucial central tropical Pacific was still normal, the CPC group came out with its first solid prediction: "It seems most likely that warmepisode conditions will develop in the tropical Pacific during the next 3-6 months." That didn't contradict the majority of model forecasts, and it fit what CPC researchers had been seeing in the changing circulation of atmosphere and ocean.

Not everyone agreed, however. "A lot of us felt they were too quick" to call for an El Niño, says meteorologist Anthony Barnston of Columbia University's International Research Institute for Climate Prediction (IRI) in Palisades, New York. On its public Web site, the IRI group started in January

> with a 60% probability of an El Niño developing and built more or less steadily to a June forecast with a 90% chance. At the same time, the models were developing a consensus for an El Niño peaking at weak-to-moderate warmth next winter. "Overall, the models are doing better this time around," says Barnston, who main-

ScienceSc⊕pe

Museum Stays Nanjing city officials have agreed to revise plans for boosting tourism to accommodate a science museum being built by the Nanjing Institute of Geology and Paleontology.

The institute, part of the Chinese Academy of Sciences, spent 4 years winning approval for a three-story, \$3.6 million museum that would display fossils and other artifacts. But in February, city officials ordered the institute to halt work on the building, to be completed next year, because it interfered with plans to enhance a nearby 1400-year-old Buddhist nunnery (Science, 24 May, p. 1379). Last month the city backed off, however, saying it will develop new beautification plans that take the museum into account.

Yang Qun, deputy director of the institute, says he's "glad that the government has reiterated its support" for the museum.

NIMH Short List The search is winding down for a new director for the National Institute of Mental Health (NIMH), which has lacked a permanent leader since Steven Hyman returned to Harvard in December 2001. A search committee has forwarded four names to National Institutes of Health (NIH) director Elias Zerhouni, sources tell Science. They are Edward Scolnick, executive vice president for science at Merck & Co. Inc.; Thomas Insel, a former NIMH researcher who is now at the Yerkes Regional Primate Research Center at Emory University in Atlanta (see p. 506); Dennis Charney, who leads NIMH's intramural program on anxiety disorders; and David J. Kupfer of the University of Pittsburgh.

The front-runner is said to be Scolnick, a former NIH cancer researcher now on NIMH's advisory council. But rejoining the government would mean a hefty pay cut.

Forgive and Take In an unusual deal, Russia has agreed to forgive \$98 million in Armenian debt in exchange for control of four state enterprises, including a pair of scientific institutes. The biggest prizes are the Hrazdan thermal power station and the Mars joint-stock company, a circuitboard manufacturer. But the inclusion of two Yerevan-based electronics research labs irks some observers. "This is a new form of neocolonialism," grouses an official of a science foundation in Yerevan. The parliaments of both countries are expected to ratify the deal this fall.

Contributors: Andrew Lawler, Jocelyn Kaiser, Ding Yimin, Richard Stone

tains a summary plot of model forecasts.[‡] "Quite a few anticipated the onset, and we had only a few real losers."

The computer models might be doing better, but they haven't won the day yet. Kousky attributes his group's success to "a combination of more experience [watching El Niños develop], 2 decades of research, and the observation network [the National Oceanic and Atmospheric Administration] and NASA have invested in. We look to the models to confirm what we're seeing in our monitoring." What they're seeing now is enough warmth in the central tropical Pacific (a 3month sea-surface temperature average 0.5°C greater than normal) to justify declaring the early stages of an El Niño. They also see only enough heat stored around the Pacific to anticipate weak-to-moderate warmth this winter. Time will tell. -RICHARD A. KERR

SUPERCONDUCTIVITY

Stripes Theory Beset By Quantum Waves

A new explanation of how high-temperature superconductors get their stripes could upend a controversial theory of how some of the materials conduct electricity without resistance. According to the "stripes" theory, electric charges within copper-and-oxygenbased superconductors collect in long lines, and pairs of charges then glide along these stripes unhindered. The scenario recently got a boost when experiments tripled the number of so-called cuprate superconductors sporting these apparent stripes (Science, 15 March, p. 1992). But now physicists at the University of California (UC), Berkeley, suggest that cuprate stripes might be an illusion. In a paper published online this week by Science (www.sciencexpress.org/cgi/content/abstract/ 1072640), they show how the stripes in one material might be a subtle effect of overlapping quantum waves of electric charge.

The material is bismuth strontium calcium copper oxide, abbreviated BSCCO and nicknamed bisco. The new results do not rule out charge stripes in BSCCO, but they show that stripes are not needed to explain the undulations earlier researchers spotted on the surface of the material, says Juan-Carlos Campuzano, a physicist at the University of Illinois, Chicago. "These experiments were touted as the proof for stripes," Campuzano says. "Well, that proof will have to come from somewhere else. One does not require stripes to explain the patterns seen in these experiments." Proponents of stripes, however, say the UC Berkeley team's data contain evidence of stripes that the experimenters themselves have overlooked.

On one issue all agree: Those data give the best look so far at BSCCO's electrical properties. To get them, J. C. Séamus Davis and his UC Berkeley colleagues studied the surface of a single crystal of BSCCO with a tiny, fingerlike electrical probe called a scanning tunneling microscope (STM). As they slowly moved the probe across the surface, the current between the tip and the material rose and fell in ripples—a pattern that other researchers had ascribed to stripes of charge lying just beneath the surface.

Going beyond the earlier studies, the UC Berkeley researchers found that as they varied the voltage between tip and surface, the spacing of the ripples changed. That shouldn't happen if stripes were causing the ripples, they argue. Instead, the spacing would stay the same regardless of the voltage —much as the spacing of the pickets in a fence feels the same no matter how hard you press when running your finger over them. Moreover, Davis and colleagues found that

the spacing varied just as it would if the ripples sprang from a quantum wave of electrical charge, reflecting off an imperfection in the crystal and then interfering with itself to produce peaks and valleys. They compared their results with data Campuzano and colleagues had taken by another technique, which showed just which wavelengths and energies such waves might have in

Bumps on the right signal stripes—if they're there.

Peaking through.

BSCCO. The two types of data agree nearly perfectly, Davis says.

Davis acknowledges that stripes might still appear under other conditions—if BSCCO were exposed to a magnetic field, say, or if its composition were changed. But quantum waves explain all the ripples the researchers see in this experiment, he says: "Ockham's razor says that this is all that's necessary."

Yet some researchers argue that the UC Berkeley group shaved away the stripes in its own data. The STM readings also show small signs of ripples whose spacing does not change with voltage, says Aharon Kapitulnik, a physicist at Stanford University in Palo Alto, California, who previously studied BSCCO with an STM and says he saw stripes. In plots where Davis and colleagues see only one peak that moves as the voltage changes, he sees a second that doesn't (see figure). "It's in their own data." Kapitulnik says.

Steven Kivelson, a physicist at the University of California, Los Angeles, thinks both waves and stripes might be playing a

role. Contrary to what Campuzano's data predict, Kivelson says, the spacing of Davis's ripples appears to stop changing as the voltage between tip and sample decreases toward zero. That suggests the wave phenomenon fades and stripes emerge at very low voltages. If such subtle signs of stripes are there, Davis and colleagues should find them as they push forward with their technique.

Meanwhile, a fresh challenge to the stripes model comes from a different direction. On page 581, Peter Abbamonte and colleagues at the University of Groningen in the Netherlands and Oxxel GmbH in Bremen, Germany, report a new x-ray scattering technique that should be especially sensitive to charge stripes. The researchers zapped a film of superconducting lanthanum copper oxide (LCO) with x-rays tuned so they'd be absorbed in regions where the LCO contained extra electrical charge. That would cause the

x-rays to reflect in a way that revealed the presence of charge stripes. The researchers saw no evidence of such an effect.

Unfortunately, the samples Abbamonte and colleagues examined are not quite like the other lanthanumbased materials that have already shown their stripes, says John Tranquada of Brookbayen National Labora

haven National Laboratory in Upton, New York. Thus it's hard to know whether there should have been stripes for the Dutch-German researchers to see. "Because it's a new technique, it would be useful to calibrate it on one of the exact same materials where we have evidence for stripes," he says. Abbamonte, now at Cornell University, says he's currently trying to do just

that by studying samples in which Tranquada spotted stripes in 1995.

Even if all cuprate superconductors have stripes, physicists still have to determine whether they really work the way the stripes theory predicts, Tranquada says. That could be an even tougher job than reconciling contradictory results and interpretations of the data, and it might keep physicists reading between the lines for some time to come.

-ADRIAN CHO

Adrian Cho is a freelance writer in Grosse Pointe Park, Michigan.

