

inoculations were successful in 50 per cent. of the attempts. The controls were negative in over 90 per cent. of the patients. These results seem to indicate that *Pityrosporum ovalis* may produce seborrheic dermatitis under favorable conditions.

MORRIS MOORE  
ROY L. KILE

THE BARNARD FREE SKIN AND  
CANCER HOSPITAL  
ST. LOUIS, MISSOURI

### A DWARF MUTATION IN THE RAT

In rodents genetic dwarfs have been reported in the guinea pig,<sup>1,2</sup> in the mouse by Snell<sup>3</sup> and recently in the rabbit by Green, Hu and Brown.<sup>4</sup> The dwarfism in the mouse was shown by Smith and MacDowell<sup>5</sup> to be caused by a hereditary deficiency of the anterior pituitary. Preliminary studies in the rabbit suggest that the defect may be due to an endocrine abnormality, while in the guinea pig the nature of the dwarfism was not determined.

The dwarf mutation in the rat described herein was first observed in our colony in the summer of 1933. It appeared as one individual in a litter of nine young in a strain of rats that had been closely inbred in our laboratory for several years. Shortly afterward another dwarf was observed in a litter from a closely related pair, and subsequently in the litters of other closely related rats. At present 22 dwarfs have been produced in 12 litters, the ratio of normal to dwarf in these litters being 80 to 22. These results suggest that the dwarf is the result of a simple autosomal recessive mutation, the appearance of both dwarf males and females showing that the gene is not sex-linked.

At birth normal and dwarf rats can not be distinguished, and it is impossible to separate the dwarf rats with certainty until about the twelfth day after birth. At this time a distinct difference in rate of growth becomes manifest, and the hair of the dwarfs appears much thinner and of finer texture than the hair on their normal litter mates. This difference in the hair is a characteristic feature of the dwarfs throughout life. The mature weight of the dwarfs is approximately 50 per cent. that of the normal males and 70 per cent. of the normal females of this strain. A remarkable feature is the failure of the males and females to become differentiated in size, whereas the male normally becomes distinctly larger than the female.

Thus far all dwarfs have proven to be sterile. No sexual activity has been observed in dwarf males and in only one instance has copulation been noted among the dwarf females. The size of the testes remains infantile, becoming less than one-half normal size. Spermatogenesis occurs, but at a greatly reduced rate, and the few sperm observed were abnormal in appearance. As yet studies have not been made of the ovary.

In all cases so far observed a distinct opacity of the lens of the eyes has been associated with the dwarfism, although some variation in the degree of this opacity exists. Skeletons of the dwarfs have not been prepared, but the general proportion of the body parts appears the same as in normal rats. The effect seems to be a general reduction in size of all parts, and the retardation in rate of growth becomes manifest at a very early date. Using the method of analysis of variance on the weights of litters in which dwarfs appeared, it has been determined that the retardation in growth had begun on the fifth day. As pointed out above, however, individual dwarfs can not be separated with certainty until about the twelfth day after birth. The dwarf rats are weaker than normals, are more susceptible to infections, and are shorter lived.

Individuals heterozygous for the dwarf gene are normal in appearance, in vitality and they attain the same size as homozygous normal rats. Recently, however, some individuals have been observed among the progeny of dwarf-producing pairs that were slightly slow in developing their hair and whose early growth seemed somewhat retarded. Tests are now under way to determine whether such individuals are heterozygous for the dwarf gene.

The decreased size of the dwarf rats, their sterility and their general appearance suggests that the defect may be of endocrine origin. It is, however, apparently not due to a pituitary deficiency, since in preliminary tests with the implantation of pituitary bodies from normal rats, following the technique of Smith and MacDowell, growth was not produced in treated dwarfs. Further experiments to determine the physiologic cause of this dwarfism are at present under way and the colony is being expanded in order to insure a constant supply of the dwarfs.

W. V. LAMBERT  
A. SCIUCHETTI

IOWA STATE COLLEGE

<sup>1</sup> I. J. B. Sollas, Repts. Evol. Com. Roy. Soc., 5: 51-79, 1909.

<sup>2</sup> I. J. B. Sollas, Jour. Gen., 3: 201-204, 1914.

<sup>3</sup> G. D. Snell, Proc. Nat. Acad. Sci., 15: 733-734, 1929.

<sup>4</sup> H. S. N. Green, C. K. Hu and W. H. Brown, SCIENCE, n. s., 79: 487-488, 1934.

<sup>5</sup> P. E. Smith and E. C. MacDowell, Anat. Rec., 46: 249-257, 1930.

### BOOKS RECEIVED

CAULLERY, MAURICE. *French Science*. Pp. xi+229. Illustrated. Author.  
EDDINGTON, SIR ARTHUR. *New Pathways in Science*. Pp. x+333. 4 plates. Macmillan. \$3.00.