

of Algonkian time and that the sediments were accumulated in fresh-water lakes of huge size situated at a considerable distance from the margins of the continents. In these lakes was relatively little animal life. This view is not in accordance with the testimony of the sediments, for Barrell⁴ has shown that many horizons of the Belt series of northern United States and southern Canada and of the Grand Canyon series strongly indicate deposition below sea level. My detailed studies at the Grand Canyon and some acquaintance with the Belt series, the Apache group of southern Arizona and the unnamed Algonkian series of southeastern California support Barrell's findings. The hypothesis originally advanced by Brooks⁵ and recently modified and expanded by Raymond⁶ seems a much more reasonable explanation of the paucity of fossils. These authorities and others hold that animals developed a lime- or silica-secreting mechanism for the generation of resistant parts relatively late in their evolution and that this had not taken place in most forms by Algonkian time. Even in the early and middle Cambrian, the secretions were mostly of chitin, but by the late Cambrian many forms developed structures of calcium carbonate and silica. Raymond holds that most pre-Cambrian animals were motile, swimming or crawling forms, lacking in hard parts. Predaceous carnivores apparently were scarce, hence, "the swimming and floating organisms must have increased rapidly until there came a time when the upper, sunlit part of the oceans was over-populated. This would force some individuals to the bottom. . . . Those animals which reached the bottom near the shore, where the waters were shallow, found abundant food, and survived. . . . Active animals reaching the bottom continued to swim, or learned to crawl after food. The more passive forms adhered to the substratum, became relatively inactive, and began the secretion of skeletons because they were no longer able to get rid of the calcium carbonate. . . . Animals had only commenced to discover the bottom in early Cambrian times." Thus it is probable that the Algonkian oceans were teeming with life when they spread over northern Arizona, but most of the animals, being composed only of soft tissue, left no record. Strange it is that a jelly-fish, one of the most perishable of animals, should have left the only imprint so far discovered. Many fossils doubtless are present in the great mass of sediments deposited in the Grand Canyon region, but so widely scattered are they that great masses of rock may be examined without finding any. Search in the Algonkian Apache beds of southern Arizona, probably

correlative with the Unkar strata at Grand Canyon, so far has not yielded any animal fossils.

In the lower Unkar and especially in the Chuar group, are many limestones which exhibit the structures commonly described as of algal origin. Some of these probably are inorganic, but many of them seem without question to represent algal secretions. The number of such horizons indicates that, at various times at least, the seas were heavily populated with these primitive plants. Most of the limestones probably are marine, but there is the possibility that some may have been accumulated in fresh-water lakes existing on the flood-plains when the land was above sea level. A detailed study of the considerable variety of these structures in the Grand Canyon Algonkian would be a material contribution to pre-Cambrian paleontology. There is no record of any higher types of plants.

Another indication of the abundance of life during Chuar (later Algonkian) time is the great volume of carbonaceous shale and the smaller quantity of carbonaceous and bituminous limestone. Examination of much of this material which I collected has not yielded any actual fossils. Some of the limestones are so strongly bituminous as to emit a very distinct fetid odor when broken.

NORMAN E. A. HINDS

UNIVERSITY OF CALIFORNIA

A RAINBOW AT NIGHT

I WOULD hesitate to record so trifling an observation as I recently made, were it not for the fact that none of my scientific friends with whom I have spoken seem to be aware of the phenomenon. On the night of June 16, 1938, I observed a rainbow caused by the moon, then only three or four days beyond its full stage. I was crossing from Nassau to Miami on the *Ena-K*, a small motor launch, and was obliged to remain on deck all night. The moon rose about 9 o'clock out of a beautiful calm sea. There was no land in sight. Tumultuous trade wind clouds towered to gigantic heights and there were occasional squalls of rain. About 11 o'clock, when the moon was well up in the southeast sky, the rainbow appeared in the northwest, where a thunderstorm was in progress. The prismatic colors were fairly distinguishable. The arc was complete, the two ends dipping into the sea. At no time was there the least doubt as to the cause of the phenomenon. The conditions were unusually favorable, but probably no more so than at every occurrence of full moon in the trade wind belt, where thunder squalls are common. June is their so-called rainy season for this reason.

A. K. LOBECK

COLUMBIA UNIVERSITY

⁴ J. Barrell, *Jour. Geol.*, 14: 553-560, 1906.

⁵ W. K. Brooks, *Jour. Geol.*, 2: 455-479, 1894.

⁶ P. E. Raymond, *Bull. Geol. Soc. America*, 46: 375-391, 1935.