

than for the other altitudes except the lowest at which samples were taken. The plates showed also numerous fungus spores, plant hairs and pollen from other species of plants, notably *Pinus* spp. One pine pollen grain collected at 4,000 feet above the valley floor had germinated on the agar.

It seems at this time desirable merely to record the presence of viable sugar-beet pollen in the air at high elevations without any inferences as to relation to cross-pollination problems of the sugar beet.

FRED C. MEIER  
ERNST ARTSCHWAGER

EXTENSION SERVICE  
BUREAU OF PLANT INDUSTRY,  
U. S. DEPARTMENT OF AGRICULTURE

### BASAL DIETS FOR VITAMIN B<sub>1</sub> DETERMINATION

INVESTIGATIONS which are still in progress have demonstrated that it is quite practical to prepare a diet satisfactory for the determination of vitamin B<sub>1</sub> by applying the observation of Williams and coworkers<sup>1</sup> that this vitamin is destroyed by cleavage with sulfite. A basal diet consisting of sucrose, 71 per cent., vitamin B<sub>1</sub> free casein, 18 per cent., salt mixture, 4 per cent., fat, 5 per cent., and cod liver oil, 2 per cent., was used, and various proportions of sucrose were replaced by addenda containing the vitamin B complex.

Sulfite treatment of yeast was carried out as follows: 400 cc of 0.1 per cent. sodium sulfite were added to 50 grams of dried yeast in a 500-cc wide mouth bottle. SO<sub>2</sub> was introduced until a pH of 4 was reached, and the bottle was then stoppered and allowed to stand 5 days at room temperature (25° C.). The contents of the bottle were then dried on purified casein at a temperature not exceeding 65° C.

Rats fed the basal ration containing 5 or 15 per cent. of sulfite-treated yeast, and receiving in addition crystalline vitamin B<sub>1</sub>, grew as rapidly as animals receiving the same quantity of untreated yeast in the basal diet. Six animals 27 days old, weighing approximately 40 grams, were fed the basal ration with 15 per cent. of sulfite-treated yeast. Four of these animals developed acute polyneuritis in 32, 33, 34 and 34 days, respectively. Two animals died in 24 and 31 days, respectively, the latter showing slight symptoms of polyneuritis before death. There is reason to believe that with slight modification of the basal diet the percentage of polyneuritis can be increased and animals may be produced which are more suitable for quantitative assay where duration of cure of polyneuritis is used as the criterion.<sup>2</sup>

<sup>1</sup> R. R. Williams, R. E. Waterman, J. C. Keresztesy and E. R. Buchman, *Jour. Am. Chem. Soc.*, 57: 536, 1935.

<sup>2</sup> O. L. Kline, C. D. Tolle and E. M. Nelson, *Jour. Assoc. Off. Agric. Chem.*, 21: 305, 1938.

In the preparation of rations which contain adequate amounts of the members of the vitamin B complex and devoid of vitamin B<sub>1</sub>, the destruction of vitamin B<sub>1</sub> with sulfite appears to offer definite advantages over any procedure that has been proposed. Details of an exact procedure are being studied, but because of a widespread interest at present in methods for determining the vitamin B<sub>1</sub> content of foods and pharmaceutical preparations as well as human requirements for vitamin B<sub>1</sub>, this preliminary report may be helpful to others.

O. L. KLINE  
CHESTER D. TOLLE  
E. M. NELSON

FOOD AND DRUG ADMINISTRATION,  
U. S. DEPARTMENT OF AGRICULTURE

### BOOKS RECEIVED

- Actualités Scientifiques et Industrielles; Nutrition*, 563, VIII, *Les Régulations Hormonales du Métabolisme Glucidique*. Pp. 92. 20 fr.; 576, IX, *Homéothermie et Thermorégulation, I, L'Homéothermie*. Pp. 70. 15 fr.; 577, II, *La Thermorégulation*. Pp. 73. 15 fr.; 581, XI, *Nutrition et Dentition*. Pp. 76. 20 fr.; 582, XII, *Le Problème du Pain, III, La Qualité des Blés et Son Amélioration*. Pp. 43. 10 fr.; 594, *Physiologie; 1, Glucides*. Pp. 66. 15 fr.; 2, *Eau et Sels*. Pp. 27. 8 fr.; 3, *Protides*. Pp. 54. 12 fr.; 4, *Substances Nucléiniques*. Pp. 30. 8 fr.; 5, *Créatine et Créatinine*. Pp. 40. 12 fr. Hermann, Paris.
- BLAIR, G. W. SCOTT. *An Introduction to Industrial Rheology*. Pp. xiii + 143. 20 figures. Blakiston.
- BREDERECK, HELLMUT and ROBERT MITTAG. *Vitamine und Hormone und Ihre Technische Darstellung; Erster Teil, Ergebnisse der Vitamin- und Hormonforschung*. Pp. xii + 138. Hirzel, Leipzig.
- CARMICHAEL, EMMETT B. *Laboratory Manual of Physiological Chemistry*. Pp. xxi + 126. University of Alabama. \$2.10.
- IMUS, HENRY A., JOHN W. M. ROTHNEY and ROBERT M. BEAR. *An Evaluation of Visual Factors in Reading*. Pp. xiv + 144. 11 figures. Dartmouth College.
- MADISON, HAROLD L. *Wild Flowers of Ohio*. Pp. 190. Illustrated. Museum of Natural History, Cleveland. \$1.50.
- Science Reports of the Tôhoku Imperial University; Fourth Series (Biology), Vol. XIII, No. 2; October, 1938*. Pp. 139. Illustrated. Maruzen, Tokyo.
- Scientific Reports of the Australasian Antarctic Expedition, 1911-14; FRED A. BAGE; Vol. II, Part 6, Crustacea Decapoda*. Pp. 13. 1 plate. 3/. Vol. II, Part 7, *Cirripedia*. Pp. 14. 4 plates. 1 figure. 3/6d.
- MAURICE BURTON. *Non-calcareous Sponges*. Pp. 22. 1 figure. 3/6d. Government Printer, Sydney, Australia.
- Scientific Reports of the John Murray Expedition, 1933-34; Vol. V, No. 4, by N. B. EALES, A Systematic and Anatomical Account of the Opisthobranchia*. Pp. 122. 1 plate. 28 figures. 5/. Vol. V, No. 7, by J. STANLEY GARDINER and PEGGY WAUGH, *The Flabellid and Turbinolid Corals*. Pp. 167-202. 7 plates. 6 figures. 5/. British Museum, London.
- STAMP, L. DUDLEY. *Physical Geography and Geology*. Pp. vii + 256. 199 figures. Longmans, Green. \$1.75.
- TRUDEL, WALTER. *Die Alamannen von Elgg, Eine anthropologische Untersuchung*. Pp. 114. 12 figures. G. Buchi, Zurich.