


AVAILABLE FOR INVESTIGATORS IN THE VITAMIN FIELD

Pyrazine-2, 3-dicarboxylic acid
Pyrazine monocarboxylic acid
2-methyl-1, 4-naphthoquinone
Phthiocol (2-methyl-3-hydroxy-
1, 4-naphthoquinone)

*These chemicals, prepared in our research laboratory, will
be supplied to qualified investigators upon application.*

MEAD JOHNSON AND COMPANY
EVANSVILLE, IND., U.S.A.



New!

Operates
continuously
without
becoming
too hot
to handle.

ADAMS SCOPELITE . . .

Attractively moulded of durable, almost unbreakable Bakelite, it is cool and efficient in operation. Can be used either as a substage lamp (with mirror removed) or in front of the microscope.

The louvers, front and rear, cause a free circulation of cooling air throughout the lamp without loss of light, so that even after extended use, the lamp can be handled. The frosted daylight filter diffuses the light and gives uniform illumination approximating daylight.





A-1493 ADAMS Scopelite each \$2.50

Write for quantity discounts

CLAY-ADAMS CO., 44 East 23rd St.
Inc. New York, N. Y.



Use
POLAROID* FILM
for Economy



In classroom and laboratory experiments

Polaroid Type II Film is in the form of durable, flexible sheets, easily cut with scissors to fit improvised apparatus for working with polarized light in projectors, microscopes, polariscopes, polarimeters, refractometers, pocket mineralogy polariscopes, densitometers, light intensity controllers, 3-dimensional picture demonstrations, photometers, and devices for scientific exhibits. It is economical. Nine pairs of microscope attachments may be cut from one standard 4" square of film—at a cost of less than 50c each. Two sheets, 4" square, price: \$7.50. Write Division 1... for catalog.

POLAROID CORPORATION

721 Main Street, Cambridge, Mass.

*T. M. REG. U. S. PAT. OFF.

Determination of Ammonia

REAGENT—Thymol

METHOD—Colorimetric

ADVANTAGES—Sensitivity . . . accuracy . . . stability of color

REFERENCE—Hansen and Nielsen, *J. Biol. Chem.*,
131, 309 (1939)

A GREENISH-BLUE COLORATION is formed by ammonia with thymol in the presence of hypobromite. Extraction of this color with iso-propyl ether gives a pink to deep red solution, the depth of color being directly proportional to the concentration of ammonia. Measurement is made in a Pulfrich photometer. The method is simple, accurate, and sensitive to small quantities of ammonia.

Eastman Thymol, No. 248, is a purified grade that is satisfactory for use in this analysis. An abstract of the procedure will be forwarded upon request. . . . Eastman Kodak Company, *Chemical Sales Division*, Rochester, N. Y.

Visit the Kodak Building at the New York World's Fair

EASTMAN ORGANIC CHEMICALS

WELCH

Announces

POOL'S NUCLEAR TRANSMUTATION CHART

A comprehensive and up-to-date picture of transitions existing between stable and radio-active nuclei of all the elements.

REACTIONS OF OVER 500 ATOMIC NUCLEI PORTRAYED ON THE CHART

280 Stable Nuclei—those of the more common elements—showing Isotopes and their relative percentages of abundance—

44 Natural Radio-active Nuclei, the best known being those of radium, radon and uranium—

246 New Artificial Radio-active Nuclei—
Various nuclear reactions produced by bombardment.

• 4 DISTINCTIVE COLORS • Size 42 x 50 inches
WITH MANUAL BY M. L. POOL—No. 4856, Each **\$5.00**

W. M. Welch Scientific Company

Established 1880

1517 Sedgwick Street

Chicago, Illinois, U.S.A.

*Write for complete
illustrated catalog*