

hence, the first possibility is that at the quake itself the energy is spread over such a wide area that the waves emitted, unless brought to a focus, possess too little energy to make a perceptible record.

The other possibility is that the seismologists receive waves only from the real image of the hypocenter and not even feeble ones from the hypocenter itself. In this case we should have to postulate that the waves from the hypocenter were screened from the observing stations. The nature of this screening brings us to the question of how such a real image could be formed.

If we adopt Airy's view of mountain roots, it is easy to picture the roots as forming a vast parabolic basin which would act as a mirror with the reflecting surface the denser material in which the mountains may be considered as floating. If we take the Himalayas or the Hindu Kush Mountains where the quakes which occur are consistently of the same depth of focus, about 200 kms, the mountain roots would have to form a parabolic mirror extending this depth into the isostatic level and in length and width extending roughly the length and width of the mountain range.

Such a parabolic mirror, formed by sufficiently deep roots of a mountain range such as the Himalayas, could conceivably account for a real image of a surface fracture such as we have pictured, but the mountain roots would have to be deeper than previously imagined. For land quakes such as those occurring in the Hindu Kush Mountains, the fracture might occur above sea level, and hence only those rays brought to a focus below would be recorded. Those traveling direct from the fracture would strike the side of the mountain and give rise to but very feeble surface waves. For sea quakes, *i.e.*, quakes with their origin in sub-oceanic areas, we should have to postulate a screening effect by internal reflection from the sides of the parabolic mirror.

In conclusion, it must be emphasized that there is no evidence so far adduced to show that such image foci actually exist, but on the other hand there seems to be no evidence to show that they can not exist. Their existence is suggested as a possible alternative to overthrowing the existing geological picture of an earth with a rigid outer structure extending only some sixty or seventy miles. One step toward testing the theory is being made. Records of deep-focus quakes are being carefully scanned for evidence of phases earlier than the regular P phases. Naturally, one would not expect perfect screening of waves from an extended fracture and if there are two such hypocenters, a real one and its image, one should expect at least occasionally to pick up waves from each.

JOSEPH LYNCH, S.J.

FORDHAM UNIVERSITY

RESISTANCE TO SULFANILYL DERIVATIVES IN VITRO AND IN VIVO

DURING the past year and a half, we have characterized the growth of bacteria in broth by two constants, the time T required to reach a slight but definite turbidity,¹ and the slope K of the logarithmic increase thereafter. Organisms grown in the presence of sulfanilamide, sulfapyridine, or sulfamethylthiazole can show a marked increase in T accompanied by a much smaller change in K , indicating the development of resistance after a period of inhibition.

This resistance is partly temporary, and partly "permanent" (*i.e.*, appears in subcultures). The "permanent" resistance may be intensified by continued subculture in the presence of the drug, as shown by others for *Pneumococcus*² and *Gonococcus*.³ We have found that eventually both T and K of the resistant strain grown in the presence of the drug approximate those of the parent strain tested in the absence of the drug.

The development of such resistance appears to be a general phenomenon. It also occurs *in vivo* during a course of therapy, as reported for *Pneumococcus* against sulfapyridine,⁴ and as we have found in *Staphylococcus aureus* against sulfamethylthiazole.

The question arises whether organisms made resistant to one drug also are resistant to related drugs. We have found that two types exist, A, those in which resistance induced by one drug (*e.g.*, sulfanilamide) is carried over, and B those in which it is not. The metabolically versatile *E. coli* and *Staphylococcus aureus* belong to type A, whereas the fastidious *Hemophilus parainfluenzae* belongs to type B.

The implications of the foregoing for the theory of drug action and the control of therapy are obvious.

JEROME S. HARRIS

HENRY I. KOHN

DUKE UNIVERSITY SCHOOL OF MEDICINE

COLLECTING SUBTROPICAL PLANTS AND ANIMALS IN NORTHERN OHIO

CONCERNING the spread of animals and plants throughout the United States, it is of interest to consider the probable roles played by the various aquatic plant nurseries, such as the William Tricker Company in Independence, Ohio, which import plants from many tropical and subtropical areas throughout the world, and, in turn, ship them to places all over the country.

During the entire year, the pools of this company serve as a reliable source of extremely interesting

¹ The optical density determined in the Evelyn photoelectric colorimeter is directly proportional to the number of organisms.

² C. M. MacLeod and G. Daddi, *Proc. Soc. Exper. Biol. and Med.*, 41: 69, 1939.

³ L. Westphal, R. L. Charles and C. M. Carpenter, *Jour. Bact.*, 39: 47, 1940.

⁴ R. W. Ross, *Lancet*, 1: 1207, 1939.