

the evidence of Theorell and Swedin^{3,4} which indicates that this new enzyme is really identical with peroxidase. We find that in the presence of fresh alcoholic dioxymaleic acid, acetate buffer of pH 4.5, partially purified horse-radish peroxidase and hydrogen peroxide, methyl red/o-carboxy-benzene-azodimethylaniline/ is rapidly oxidized and decolorized. In the absence of dioxymaleic acid the oxidation is slow. Moreover, peroxide need not be added, provided that the solution of dioxymaleic acid, methyl red, acetate buffer and peroxidase is shaken with air. In nitrogen there is no decolorization.

We find that peroxidase, as shown by testing with guaiacol and hydrogen peroxide, is rather rapidly inactivated by buffered dioxymaleic acid, and that this inactivation is retarded by aeration. Inactivated peroxidase is less effective in oxidizing dioxymaleic acid and in decolorizing methyl red than is active peroxidase.

Old alcoholic solutions of dioxymaleic acid give different results from fresh solutions. Here, a mixture of dioxymaleic acid, acetate buffer and peroxidase decolorizes methyl red even in nitrogen, while aeration largely prevents decolorization.

Our explanation of the rapid bleaching of methyl red in the presence of fresh dioxymaleic acid, peroxidase and hydrogen peroxide is that the peroxidase and peroxide convert the dioxymaleic acid into diketosuccinic acid and that the methyl red is oxidized by a coupled reaction. If one shakes dioxymaleic acid solutions in air, it is not necessary to add peroxide, since dioxymaleic acid is spontaneously oxidized and forms peroxide. This is in agreement with the evidence of Theorell and Swedin. Old solutions of dioxymaleic acid already contain peroxide, so that they require no shaking with air. We are not able to tell why shaking with air should retard the decolorization of methyl red in the presence of old dioxymaleic acid.

JAMES B. SUMNER
ALEXANDER L. DOUNCE

CORNELL UNIVERSITY

THE EFFECT OF GROWTH SUBSTANCES ON THE ROOTING OF BLUEBERRY CUTTINGS

THREE experiments have been conducted to study the effect of growth substance on the rooting of summer cuttings of blueberries, *V. corymbosum*. These experiments were carried on in 1937 and 1938, using in all over 2,500 cuttings from eleven different varieties of blueberries. Indole-3-acetic acid and indole-3-propionic acid were each used dry and in solutions of 5 and 10 mg per liter. Phenyl acetic acid was used

only in solutions of 10, 25 and 50 mg per liter. Auxilin was used at the recommended concentration No. 3. Phenyl acetic acid was the only one of the growth substances used which significantly increased the percentage of rooting and the greatest increase was at the medium concentration, 25 mg per liter. The results obtained with auxilin were very poor at the concentration used. The results of the above experiments are in agreement with those of Stanley Johnston's investigations independently conducted at the same time and reported in *Michigan Station Quarterly Bulletin*, 21: 255-8, 1939. From the results thus far obtained it does not seem advisable to recommend the use of growth substance for the rooting of blueberry cuttings.

F. B. CHANDLER
I. C. MASON

UNIVERSITY OF MAINE

THE IDENTITY OF THE TREE "ANNEDDA"

FROM your review of "The Englishman's Food"¹ it appears that the authors think the tree (called by the Indians "Annedda") which cured Jacques Cartier's men of scurvy when they were wintering at Stadacona in the winter of 1535-36, was *Sassafras officinale*. The sassafras does not grow anywhere in the Province of Quebec. Its only station in Canada is a relatively narrow strip in southern Ontario and there is no reason to suppose that it ever ranged farther north.

In support of the sassafras as against an evergreen, the authors state, according to the reviewer, that Cartier's notes particularly refer to the fact that the Indians had to wait for the leaves to appear in the spring. Perhaps authority is given for this statement, but no such passage occurs in Biggar's edition of "The Voyages of Jacques Cartier." Furthermore, Cartier says² that it was while he was walking on the ice that the Indians told him of the tree which would cure the sickness and two squaws went with him to gather some of it. Nine or ten branches were brought back, and Cartier adds: "They showed us how to grind the bark and leaves and to boil the whole in water."

The identity of the tree Annedda has been much disputed, but from considerations not necessary to discuss here, it seems likely that it was the hemlock, *Tsuga canadensis*.

CHARLES MACNAMARA
ARNPRIOR, ONTARIO, CANADA

THE KIT FOX

IN the summer of 1911, T. A. Rocklund and I were camped for one month on the Pennell Ranch in Wallace County, Kansas, adjoining the George A. Allman

³ H. Theorell and B. Swedin, *Naturwissenschaften*, 27: 95, 1939.

⁴ B. Swedin and H. Theorell, *Nature*, 145: 71, 1940.

¹ SCIENCE, 91: 217, 1940.

² H. P. Biggar, "The Voyages of Jacques Cartier," p. 212-215. Ottawa: The King's Printer, 1924.