

not only pyocyanine but all the other antibiotics tested were observed for reactivity with thiols *in vitro*. *In vivo* tests (antibacterial action) were used only as an analytical tool in measuring disappearance of antibiotic activity. We were primarily interested in inactivation as a measure of chemical reactivity.

The relevance of thioglycolate stimulation of penicillinase action or of the rapid excretion of penicillin to the mechanism of action of penicillin is nebulous at present and is certainly not evidence for or against a sulfhydryl inhibition mechanism. We do not agree that thiol inactivation of penicillin cannot proceed in broth or other such media provided sufficient inactivator is used; such media may, however, require more inactivator.

We agree with Leonard that it has not been proven that penicillin binds protein —SH groups, and it is this lack of proof which leads us to call —SH inhibition by penicillin only a theory. This author disagrees, however, with the statement that penicillin needs to bind protein —SH in direct proportion to the number of —SH groups in proteins, since we have shown that some types of —SH groups react only sluggishly or not at all with penicillin. It may be this very selectivity of the —SH reactions which endows penicillin with specificity of inhibitory action.

References

1. BARRON, E. S. G., and SINGER, THOMAS P. *J. biol. Chem.*, 1945, 157, 221, 241.
2. CAVALLITO, CHESTER J. *J. biol. Chem.*, 1946, 164, 29.
3. LEONARD, C. S. *Science*, 1946, 104, 501.

Leptospiral Infection Among Rodents in Micronesia

JOSEPH E. ALICATA

University of Hawaii, Honolulu, T. H.

Murine leptospirosis is a problem of public health importance because the causative organism, *Leptospira icterohemorrhagiae*, is capable of infecting humans and producing Weil's disease. Leptospirosis is gradually gaining recognition as a common disease of man giving rise to, among other things, jaundice, muscular tenderness, or comparatively mild symptoms. It is believed by several workers (3, 4, 5) that this disease is by no means rare, and the infrequency with which it has been reported is probably due to failure of recognition.

Although many surveys have been conducted on the incidence of leptospirosis in various parts of the world, notably Australia, Europe, India, Japan, and the continental United States, very little is known of the incidence of murine leptospirosis in the Pacific Islands. In recent years it has been shown (1, 2) that murine leptospirosis is present in the Hawaiian Islands, especially in areas having high rainfall and an abundance of rodents. Awareness of this fact indicated the need for further study of this disease in other islands of the Pacific. During July and August 1946, a preliminary survey

was conducted by the writer under the sponsorship of the University of Hawaii Pacific Islands Research Committee to determine, among other things, the possible occurrence of murine leptospirosis on the island of Moen of Truk Atoll and on the island of Ponape, both of the eastern Carolines. Of 22 rats trapped on Moen and 18 trapped on Ponape, 3 and 2,




FIG. 1. Sections of kidney of rat from Truk showing leptospirae in the urinary tubules.

respectively, showed presence of leptospirae in the urinary tubules (Fig. 1). The diagnosis was based on examination of kidney tissues fixed in formalin and stained by the silver impregnation technique (2).

As far as is known to the writer, this report represents the first record of murine leptospirosis in any of the islands of Micronesia and lends support to the belief that Weil's disease occurs in man in that area. Mumford and Mohr (6), in summarizing various diseases of man recorded from the Japanese mandated islands, state that according to the 1930 Report of the League of Nations, two cases of Weil's disease occurred in 1929 among Japanese in those islands. They also mentioned the possibility that the above cases might have been those of "infective hepatitis," and, conversely, cases which had been reported previously by other workers in the Marshalls as "catarrhal icterus" might have been cases of Weil's disease.

Although further research studies in Micronesia are necessary, the present finding points out that human leptospirosis is probably present in the area and more common than heretofore believed. The combination of high rainfall and abundance of rodents in Micronesia presents factors which are very favorable to the spread of the disease.

References

1. ALICATA, J. E. *Plantation Hlth* (Aiea, Oahu, Hawaii), 1944, 8.
2. ALICATA, J. E., and BREAKS, V. *Hawaii med. J.*, 1943, 2, 137.
3. BRUNO, F. E., WILEN, C. J. W., and SNAVELY, J. R. *J. Amer. med. Ass.*, 1943, 123, 519.
4. CILENTO, R. W. *Austr. Sugar J.*, 1935, 26, 573.
5. LARSON, C. L. *Publ. Hlth Rep.*, 1941, 56, 1593.
6. MUMFORD, E. P., and MOHR, J. L. *Amer. J. trop. Med.*, 1943, 23, 381.