

stration of a relationship between gross protein binding of hydrocarbons and a metabolic route that is specific to the carcinogenic process (12). In this study of contact carcinogenesis, there has been no identification of a specific metabolic route distinguishable from detoxication. The degradation of a hydrocarbon applied to the skin should involve enzymes, in view of which the results are reasonable. It is of interest that two examples of high metabolic activity in the skin have been found.

HERBERT I. HADLER
VIOLET DARCHUN
KATHARINE LEE

Division of Oncology, Chicago
Medical School, Chicago, Illinois

References and Notes

1. E. C. Miller and J. A. Miller, *Cancer Research* 7, 468 (1947); 12, 547 (1952).
2. E. C. Miller et al., *ibid.* 9, 336 (1949); E. C. Miller and J. A. Miller, *J. Natl. Cancer Inst.* 15, 1571 (1955).
3. E. C. Miller, *Cancer Research* 11, 100 (1951); C. Heidelberger and S. M. Weiss, *Cancer Research* 11, 885 (1951).
4. W. G. Wiest and C. Heidelberger, *ibid.* 13, 250 (1953).
5. M. M. Moodie, C. Reid, C. A. Wallick, *ibid.* 14, 367 (1954).
6. D. L. Woodhouse, *Brit. J. Cancer* 8, 346 (1954); 9, 418 (1955).
7. C. Heidelberger and M. J. Moldenhauer, *Proc. Am. Assoc. Cancer Research* 2, 24 (1955); *Cancer Research* 16, 442 (1956).
8. V. Darchun and H. I. Hadler, *Cancer Research* 16, 316 (1956).
9. This research was supported by grant C-2399 and cancer control grant CS-9212 from the National Cancer Institute, National Institutes of Health, U.S. Public Health Service.
10. H. I. Hadler et al., *Proc. Am. Assoc. Cancer Research* 2, 113 (1956).
11. P. E. Steiner and J. H. Edgcomb, *Cancer Research* 12, 657 (1952).
12. The experiments indicating the nonbinding of anthracene (5) and phenanthrene (7), which are both noncarcinogenic hydrocarbons with three rings, were not designed to detect a transient protein-bound phase.

9 July 1956

Sex Difference in Eosinophil Counts in Tail Blood of Mature B₁ Mice

Eosinophil counts in tail blood of mice vary with the phase of the daily adrenal cycle (1-4) in males (1, 4, 5) and in females (6) of several inbred strains and their hybrids. Strain differences in characteristic eosinophil levels—obtained at the daily time of high counts—have also been described (7). The study reported here (8) led to the additional detection of sex differences in the eosinophil count of mature mice of an inbred strain.

The mice studied were of a subline of the C₅₇-black stock (B₁) which had been maintained in the Division of Cancer Biology at the University of Minnesota by brother-to-sister mating for more than 20 generations. The animals were 8¼ months ± 1 week of age at the time of study. For 7 days prior to the study, they were singly housed in a room maintained

at 25.6 ± 0.6°C and illuminated from 6 A.M. to 6 P.M. and darkened from 6 P.M. to 6 A.M. Purina Fox Chow and tap water were available to the mice from the time of weaning and throughout the study. Except for the activities associated with cleaning the cages, feeding the mice, and filling the water bottles, the animals were not intentionally disturbed from birth until the time of study. The study was begun at 8:30 A.M. and ended at 8:30 A.M. the following day. It involved eosinophil counts on tail blood obtained from separate groups of mice at 4-hour intervals. Each mouse was thus used for venesection only once. The assembly-line procedures employed for eosinophil counts have been described (7).

Figure 1 shows the mean count ± 1 standard error for the two sexes. The 24-hour eosinophil rhythm stands out clearly for each sex, in agreement with the results of earlier work on this subject. But the level around which the mean count cycles in the two sexes, which was previously not compared, is not the same in the two sexes as far as the stock and age-group studied are concerned (Fig. 1). The females exhibit lower counts than the males, without any overlap of mean counts throughout the 24-hour period. Subsequent work on the same mice revealed that the mean paired adrenal weights of the females were higher than those of the males and that the ascorbic acid concentrations in glands from females were higher than in glands from males.

The observation of a sex difference in adrenal weight of B₁ mice extends to this stock observations on the sexual dimorphism of the adrenals, which was reported earlier for several animal forms (9). The sex difference in eosinophil count deserves more study, for it may be related to sex differences in adrenal secretory behavior, as anticipated but not yet reliably established (9).

The possible relation of the sex difference in eosinophil level to a sex difference in adrenal secretory activity comes to mind in view of the known eosino-




Fig. 1. Sex difference in mean blood eosinophil counts of B₁ mice approximately 8 months old. The number of mice used given above each point; serially independent sampling.

penic effect of both cortical and medullary adrenal hormones (2). The role played by interaction of the longer estrus cycle (10) with the daily adrenal cycle (2, 4, 6) which maintains the eosinophil rhythm must also be considered. But whatever may underlie the differences noted, it seems fair to conclude that, in B₁ mice of the age group studied, eosinophil counts can be shown to be a function not only of strain and of phase of daily cycle but also of sex.

FRANZ HALBERG
OLGA HAMERSTON
JOHN J. BITTNER

Division of Cancer Biology, Department of Physiology, University of Minnesota Medical School, Minneapolis, and Cambridge State School and Hospital, Cambridge, Minnesota

References and Notes

1. F. Halberg and M. B. Visscher, *Proc. Soc. Exptl. Biol. Med.* 75, 846 (1950).
2. F. Halberg, *J. Lancet* 73, 20 (1953).
3. ——— and M. B. Visscher, *Endocrinology* 51, 329 (1952); F. Halberg, *Acta Med. Scand. Suppl.* 302, 117 (1955).
4. H. E. Brown and T. F. Dougherty, *Endocrinology* 58, 365 (1956).
5. H. Panzenhagen and R. Speirs, *Blood* 8, 536 (1953); C. Louch, R. K. Meyer, J. T. Emlen, *Proc. Soc. Exptl. Biol. Med.* 82, 668 (1953).
6. F. Halberg, M. B. Visscher, J. J. Bittner, *Am. J. Physiol.* 174, 109 (1953).
7. F. Halberg, J. J. Bittner, M. B. Visscher, *Blood* 6, 832 (1951).
8. This investigation was aided by the Elsa U. Pardee Foundation and by research grants from the U.S. Public Health Service (G-4158 and B-980), the American Cancer Society, and the Graduate School of the University of Minnesota.
9. F. A. Hartman and K. A. Brownell, *The Adrenal Gland* (Lea and Febiger, Philadelphia, 1949), p. 325; R. Bachmann, in *Handbuch der mikroskopischen Anatomie des Menschen* (Springer, Berlin, 1954), vol. 6, p. 952.
10. G. Bourne and S. Zuckermann, *J. Endocrinol.* 2, 268 (1940).

29 October 1956

Tentative Correlation of Alaskan Glacial Sequences, 1956

A tentative correlation of the Cook Inlet and other Alaskan glacial sequences was published in 1953 (1). Since then, additional field and radiocarbon data have accumulated that necessitate some revisions in the 1953 correlation chart. These revisions (Fig. 1) include additions to earlier published glacial sequences and changes in correlation with the type Pleistocene chronology.

Correlations with the standard North American Pleistocene chronology have been revised in accordance with new radiocarbon data relating to both the Mid-continent and Alaskan sequences. Runs of radiocarbon samples W-76, W-77 (2), and also W-174 (3) by the more accurate gas-counting method (these samples were originally analyzed by the solid-carbon counting method) indicate that the lower boundary of the