

parisons of gamma globulin prepared from the same plasma are needed to establish this.

Gaines and Landy (5), employing a hemagglutinin technique, demonstrated antibodies in normal human serums against a *Pseudomonas* antigen; higher titers were present in older age groups. Various investigators have demonstrated small amounts of *Staphylococcus* antitoxin in the serums of man and animals, and the antibacterial properties of normal serum have been widely investigated for many years (3). The role of these *in vitro* properties of serum in its protective action against lethal infections in animals remains to be established. However, Cameron (6) has demonstrated protective action from gamma globulin derived from several domestic animals against experimental infections with *Pasteurella multocida*, *Salmonella choleraesuis* and *Brucella suis*.

R. CARL MILLICAN
J. RUST

S. M. ROSENTHAL

National Institute of Arthritis and
Metabolic Diseases, National Institutes
of Health, Bethesda, Maryland

References and Notes

1. S. M. Rosenthal, R. C. Millican, J. Rust, *Proc. Soc. Exptl. Biol. Med.* 94, 214 (1957).
2. We wish to acknowledge the cooperation of the American Red Cross for supplying the gamma globulin used in this study and of E. Verder and M. Landy of the National Institutes of Health, A. K. Miller of the Merck Institute, and E. Grunberg of Hoffman-La Roche, Inc., for cultures of organisms.
3. Topley and Wilson, *Principles of Bacteriology and Immunity* (Williams and Wilkins, Baltimore, Md., 1955).
4. R. C. Millican, J. Rust, S. M. Rosenthal, in preparation.
5. S. Gaines and M. Landy, *J. Bacteriol.* 69, 628 (1955).
6. H. S. Cameron, *Am. J. Vet. Research* 10, 265 (1949).

22 April 1957

Interrelationship between Certain Bacteria and the Rumen Ciliate *Dasytricha ruminantium*

The cattle and sheep ciliates of the genera *Isotricha* and *Dasytricha* have been shown to possess considerable fermentation ability (1, 2). Soluble carbohydrates are rapidly utilized by the protozoans, with the production of acids and gas and with concomitant deposition of food reserve amylopectin. The nature of the nitrogen source for the holotrich ciliates has been elusive (3). The holotrich protozoans do not easily lend themselves to microscopic observation of bacterial feeding. In well-fed ciliates the entire organism is loaded with reserve food granules, making it difficult to detect either the presence of bacteria

within the ciliates or bacteria actually being ingested by the protozoans.

Growth experiments with the holotrich protozoan *Isotricha intestinalis* have been reported (2). The ciliates grew during a short period in an anaerobic medium containing ground alfalfa, wheat, and untreated rumen fluid in inorganic saline. Counts of the protozoans during this time indicated division every 48 hours. The smaller, more abundant holotrich found in the rumen, *Dasytricha ruminantium*, did not survive in these experiments, indicating a difference in cultural requirements. It was of interest to find what these differences were and also what factors would stimulate growth of *D. ruminantium* in laboratory cultures.

Starvation of *D. ruminantium* for 48 to 72 hours decreased the number of amylopectin reserve granules and facilitated the search for bacterial ingestion by the protozoans. The starved ciliates were placed on a slide with a small amount of 0.5 percent saline and a mixture of bacteria obtained from fresh rumen fluid by centrifugation. A cover glass, ringed with Vaseline, excluded air from the entire suspension. With this arrangement the protozoans remained active and could be observed for several hours, with the use of a phase microscope. *Dasytricha ruminantium* selected and ingested small cocci, 0.5 to 0.8 μ in diameter, from the mixture. The bacteria agglutinated at the mouth located at the posterior of the organism and were gradually ingested. Occasionally, a ciliate was observed which swam about trailing many agglutinated bacteria. Small rod-shaped bacteria, 1.0 \times 1.5 μ in size, were seen also within food vacuoles in the ciliates. In several of the experiments, exposure of the bacterial suspension to triphenyltetrazolium chloride allowed more distinct observation of bacterial ingestion by the protozoans. Bacteria which stained red as a result of the deposition of the insoluble formazan were seen localized near the mouth and in the interior of the ciliate in food vacuoles.

Several bacterial strains were isolated from rumen contents in order to test their growth-promoting effect on *Dasytricha*. Following the initial isolation on a starch-alfalfa extract agar, the bacteria were grown in liquid medium of the same type and used in the ciliate cultures. Small vessels with a flat upper surface and an opening fitted with a rubber stopper allowed observation of the protozoans without exposure to air. With the aid of a dissecting microscope, the response and fate of the ciliates in each experiment could be followed easily. With cooling of the cultures from 39°C to room temperature, most of the protozoans decreased their swimming activity

and settled to the bottom of the vessel. The stale supernate was then removed with a capillary pipette, while anaerobic conditions were preserved by passing carbon dioxide into the vessel opening. Fresh 0.5 percent buffered saline containing substrate and bacteria was added every 24 hours. Streptomycin and penicillin, plus washing techniques, were used to free *Dasytricha* of most of the associated rumen bacteria.

Dasytricha ruminantium was cultured for 2 weeks in the presence of two species of rumen bacteria and the following components, given in percentage: 0.05 ground alfalfa, 0.005 cornstarch, 0.5 autoclaved rumen fluid, 0.005 glucose, and a mixture of ten vitamins. Cysteine hydrochloride, 0.01 percent, was added to insure reduced conditions. Numerous dividing ciliates were observed each day. Parallel control cultures which omitted the bacteria were dead at 96 hours. The bacterial strains which stimulated growth of *Dasytricha* were a small coccus, 0.8 μ in diameter, having a large mucoid colony, and a rod 1.0 \times 1.5 μ in size, possessing a filamentous type colony. The larger holotrich ciliates of the genus *Isotricha* were also observed feeding on rod-shaped bacteria (4). It is concluded that the holotrich rumen ciliates derive at least some of their nitrogenous requirements from the ingestion of associated bacteria.

JOSE GUTIERREZ
*Animal Husbandry Research Division,
Agricultural Research Center, USDA,
Beltsville, Maryland*

R. E. HUNGATE
*Department of Bacteriology, University
of California, Davis*

References and Notes

1. P. J. Heald and A. E. Oxford, *Biochem. J. (London)* 53, 506 (1953).
2. J. Gutierrez, *ibid.* 60, 516 (1955).
3. A. E. Oxford, *Exptl. Parasitol.* 4, 569 (1955).
4. A detailed paper is in preparation.

8 July 1957

Rapid Chemical Changes in Reconstituted Dry Milk

Dry milk is usually reconstituted into an aqueous system before use as a food, and many of the laboratory analyses of dry milk involve reconstitution with water as a step in sample preparation. Recent study of certain chemical properties of reconstituted nonfat dry milk indicated a variability in analytic results which was correlated with the age of the reconstituted samples. The purpose of this report is to illustrate some rapid chemical changes which may occur in freshly reconstituted dry milk in order to bring the phenomenon of change to the attention of those interested in the