

WILD* M-20


with incident light attachment

One of the most important advances in microscopy for research and scientific exploration is now within easy reach of the owner of a Wild M-20.

Here, in a superbly Swiss crafted instrument, capable of the finest results in all types of microscopy, is now added a new, first order capability:

With Incident Light Attachment, the M-20 permits observation and photomicrography under bright and dark field conditions, with polarization and fluorescence. Optical quality and handling convenience are fully comparable to those found in specially designed incident light microscopes.

For full information about this most versatile microscope...and its many attachments, write for Booklet M-20.



*The FIRST name in Surveying Instruments, Photogrammetric Equipment and Microscopes.

WILD
HEERBRUGG

Full Factory
Services

INSTRUMENTS, INC.

Main at Covert Street • Port Washington, New York
Port Washington 7-4843

In Canada

Wild of Canada Ltd., 157 MacLaren St., Ottawa, Ontario

Letters

Trajectory of Lunik III

Using available tracking data released by Tass, the General Electric Company has computed the trajectory of the Russian automatic interplanetary station (AIS), otherwise known as Lunik III, which obtained the first pictures of the far side of the moon.

The results showed that the 7th perigee passage of the vehicle occurred on 21 January at 12.87 hours (universal time) at a distance of 18,225 kilometers from the earth's center, and predicted entry into the earth's atmosphere on 8 March at 5.19 hours.

The U.S.S.R. had predicted 7th perigee passage on 22 January at 9.03 hours at a distance of 18,486 kilometers, and final entry into the earth's atmosphere late in March. Subsequent corrections to the initial conditions of the General Electric program were made to permit agreement with the 7th perigee time predicted by the U.S.S.R.; computations made on this basis show little change in the value for perigee distance. However, these computations place entry into earth's atmosphere in April or later.

The calculations are obviously very sensitive to initial conditions; this is due primarily to a second close approach to the moon (of about 50,000 kilometers), on 24 January. Additional tracking or sighting data are therefore needed to confirm, or permit corrections to be made in, the trajectory predictions if a meaningful entry watch is to be established. We would be pleased to receive such additional information and would undertake to rerun the computations and to advise both *Science* and others interested in the results.

J. E. MICHAELS

Space Sciences Laboratory,
General Electric Company,
Philadelphia, Pennsylvania

Evapotranspiration

It is surprising that no comments or criticisms have appeared in reference to the report by Holdridge [*Science* 130, 572 (4 Sept. 1959)] concerning a "Simple method for determining potential evapotranspiration from temperature data."

The formula

$$\text{Potential evapotranspiration (in mm)} = \left[58.93 \left(\frac{\text{Unit period of time}}{\text{No. of units of time in 1 yr}} \right) \right] \times \left[\text{Comparative plant growth mean temperature (}^{\circ}\text{C)} \right]$$

is not consistent with Holdridge's statement that "the potential evapotranspiration at a given temperature decreases proportionately along the gradient of increasing precipitation from arid to wet areas. . . ." It is difficult to see a theoretical basis for the formula.

Work by Penman [*Proc. Roy. Soc. (London)* A193, 120 (1948)], Thornthwaite [*Geograph. Rev.* 38, 55 (1948)], and Blaney-Criddle ["Water," *Yearbook Agr. (U.S. Dept. Agr.)* (1955)] indicate that such a formula, based on temperature alone, is of doubtful validity. Ramage [*Pacific Sci.* 13, 1 (1959)] found that both the Penman and Thornthwaite formulas gave values of potential evapotranspiration which were too high during the wet summer months at Hong Kong. Use of the Holdridge formula on the Hong Kong data shows the same tendency. In fact, values for potential evapotranspiration for Hong Kong for 1951-56 computed by the Holdridge formula are intermediate between those computed by the Penman and the Thornthwaite equations, respectively, and are considerably higher than observed values for an evapotranspiration battery.

Potential evapotranspiration is dependent upon a number of meteorological factors which fluctuate from day to day and from season to season, so that a formula based on temperature alone will be valid only under very restricted conditions of insulation, humidity, and wind.

TERRELL L. NOFFSINGER

Land Study Bureau,
University of Hawaii, Honolulu

Regarding the comments of Noffsinger, which I appreciate because they offer me an opportunity to clarify certain points in my report in *Science*, the following discussion is submitted.

Noffsinger's first paragraph looks convincing only because he has misquoted from my report, using in his letter "the potential evapotranspiration at a given temperature decreases. . . ." rather than, as appeared in the article, "the potential evapotranspiration rate at a given temperature decreases. . . ." Naturally, potential evapotranspiration is quite distinct from the potential evapotranspiration rate, which is equal to the mean potential evapotranspiration in millimeters per year divided by the mean precipitation in millimeters per year.

As for his statements on temperature, my formula uses the comparative plant growth mean temperature, preferably called "mean biotemperature," which discards temperatures below 0°C. The mean biotemperature is derived by summing up the positive time-unit means of temperature and dividing this total by the number of time units in the period. The workers cited, who found