tion of clinical drug evaluation is beyond the capacity of the investigator, the university, the government, and the pharmaceutical industry. A National Institute of Pharmacology with legal and scientific responsibility is essential. This would be a federally sponsored institute which would stimulate and supervise basic and clinical drug research with an emphasis on new drug investigation. The primary involvement of the FDA with food, cosmetics, manufacturing, and advertising indicates that new drug investigation should be in a separate program patterned on the National Institutes of Health. The work of such an Institute of Pharmacology should be conducted by universities and research facilities which conform to the highest standards of personnel, equipment, and research design. The pharmaceutical industry would not be relieved of its obligation to demonstrate the effectiveness and safety of its products and to underwrite the cost of this work but there would be a federal capability which would set standards and enforce them. Such a program would insure the badly needed financial support of new drug research. It would also require complete and prompt reports of new drugs which would be available to the investigators as well as to the government.

PAUL LOWINGER

School of Medicine, Wayne State University, 951 East Lafayette, Detroit, Michigan 48207

Reference

 I. T. Borda, D. Slone, H. Jick, J. Amer. Med. Ass. 205, 645 (1968).

Overhead Costs during Austerity

The austerity program for scientific research is requiring some adaptations. For example, in our department the cost of publication page charges for a 12-man faculty was \$15,000 last year. This means that page charges cost as much as an additional faculty member. We have been wondering whether the actual scientific communication achieved by the present method is worth the cost. Because we are skeptical, we are trying the following method. Work which is supplementary to an existing key publication will not be published per se but will be written up with no regard to saving space, then will be multilithed, and made available as a numbered "Supplemental Publication"

Bausch & Lomb Breaks the Cost Barrier to Laser Science

Now, every high school and college physics department can have a dependable helium-neon gas laser... to demonstrate optical principles, such as interference and diffraction patterns and to perform experiments in making and reconstructing holograms.


Teaching


This revolutionary, low-cost laser is designed to be used routinely in classroom experiments. Operation is simple and dependable, Just plug it in and it begins to lase. Its .1mw power output is ideal for classroom use. Produces a monochromatic beam at 6328 Angstroms. Power requirement is standard 115-v, AC, 50-60 cycles.

Ask for catalog 41-2306. Or better still, place your order now. Bausch & Lomb, Scientific Instrument Division, 87816 Bausch Street, Rochester, New York 14602.



Circle No. 33 on Readers' Service Card





Precision molded and precision calibrated, the Nalgene® Volumetric Flask is in a class by itself. It won't etch, contaminate or break. Repeated autoclaving won't affect its accuracy. Each flask is individually calibrated to better than ± 1/6 of 1%.

100 ml size now in stock, 250, 500, and 1000 ml sizes coming soon. Order from your lab supply dealer . . . and specify Nalgene Labware. Ask for our 1968 Catalog or write Dept. 2104, Nalgene Labware Division, Rochester, N.Y. 14602.

NALGENE LABWARE DIVISION

VISIT OUR BOOTH AT THE FASEB CONVENTION

of our department. We intend to deposit two bound volumes of these in the university library each year so that Xerox copies can be sent to anyone for the normal cost. In this way we can provide the information whenever it is needed and we can publish the key parts of the research with more brevity. We would also welcome two new avenues of publication: 1000-word summaries and current reviews covering about 10 to 15 papers in one area. We feel this latter exercise would serve to advance the faculty, rather than produce an inchoate mass of publications whose usefulness is in question. This threefold method of publication would save costs, reduce the mass of literature on the shelves, and appreciably increase scientific communication.

Our second area of adaptation causes more concern. It has been our practice to employ undergraduates and high school students in the laboratory. Many of these have become first-rate scientists: all have benefited in increased maturity and responsibility. Now we have to cut them off. If funds from an appropriate agency, such as the Office of Education, could be added to each research grant for this purpose, a real human value would be retained.

In the atmosphere of austerity the overhead which is incurred becomes of real importance and forms a strong point of division between scientists in the laboratory and administrators. Faiman's recent remarks (Letters, 27 Dec., 1968) are pertinent. I suggest that granting agencies require that the use of overhead be explained by the institution to the principal investigators, that discussion of the manner of its use be permitted, and that the agency be aware of what takes place in such discussion.

ERNEST C. POLLARD Department of Biophysics, Pennsylvania State University, University Park 16802

What Makes Oysters Grow?

A letter from Adler (29 Nov.) comments on Bardach's article "Aquaculture" (13 Sept., p. 1098). It is Adler's opinion that Bardach created a wrong impression in a statement to the effect that oyster larvae need flagellate algae for food. Adler says further that "Good larval growth has even been achieved with some nonliving substitute food like corn flour." We are not familiar with

any success at rearing larvae with nonliving foods. Either Adler has created a wrong impression of the state of marine animal husbandry, or he knows of a very important breakthrough not generally known to practitioners and scientists in aquaculture. . . .

> RICHARD J. BENOIT EDWARD A. ZURAW DONALD E. LEONE

Marine Sciences Research and Development, General Dynamics, Groton, Connecticut 06340

The search for artificial foods for marine bivalves and their larvae began in the early 1930's (1). In 1940 Bruce et al. (2) showed that live, naked flagellates were superior to some other algae for rearing larvae and spat. These studies have been confirmed many times during the past 28 years and laid the foundations for present-day hatchery techniques. Investigators have continued to search for other algae or substances which would give better results. Walne (3), evaluating the food value of seven algal species for oyster larvae, found that the diatom Phaeodactylum tricornutum promoted growth comparable to that of flagellates, while Imaya (4) indicated that both Cyclotella nana and Chaetoceros spp. are well utilized. Somewhat questionable results were obtained by Loosanoff and his co-workers (5) by feeding dried powdered Ulva and Laminaria to larvae, but when freeze-dried Schedesmas obliquus was used, Hidu and Ukeles (6) reported excellent results. Corn starch and corn meal or both were employed in oyster feeding experiments by Haven and an enthusiastic evaluation of its successful application was published by Ingle (7). Benoit and his co-workers might discover additional references by consulting the literature or the researchers mentioned above.

CYRUS ADLER

Offshore/Sea Development Corporation, 99 Nassau Street, New York 10038

References

- H. A. Cole, Fish Invest. London, ser. 2, 15, 4 (1937).
- J. R. Bruce, M. Knight, M. Parke, J. Mar. Biol. Ass. U.K. 24, 337 (1940).
 P. R. Walne, ibid. 43, 767 (1963).

- P. R. Waine, ibid. 43, 767 (1963).
 H. Imaya, personal communication; G. C. Matthiessen and R. C. Toner, Mar. Res. Found. Inc., Martha's Vineyard (1966), p. 136.
 V. L. Loosanoff and H. C. Davis, Commer. Fish. Rev. 25 (1), 1 (1963); V. L. Loosanoff, Trans. 29th N. Amer. Wildlife Natur. Resour. Conf. (1964), p. 332.
 H. Hidu and R. Ukeles, Proc. Nat. Shellfish. Ass. 53, 85 (1962)
- Ass. 53, 85 (1962).
- 7. R. M. Ingle, Sea Front. 13 (5), 296 (1967).