

have given at least some idea of what can be done with the use of such estimators, and now the time seems to be ripe for exploiting various possibilities by a large-scale systematic theoretical-numerical study. This volume contains the results of a large-scale Monte Carlo study of exactly that kind, which was undertaken at Princeton in 1970–71 under J. W. Tukey, with the participation also of P. Huber, J. Bickel, and others. Some 70 estimators are tested, including all four of the kinds described above, as well as some special ones such as the maximum likelihood estimator for the Cauchy distribution. The sampling distributions of those estimators are calculated for several shapes of distributions ranging from the normal to the Cauchy for the sample size $n = 5$ to 40.

It is difficult to assess exactly such a large quantity of numerical results, and still harder to explain them in few sentences. Moreover, the judgments of the authors themselves do not seem to be always unanimous. The general impression does not seem to be radically different from what has already been observed unsystematically, but it seems that Hampel's estimator, which is the M-estimator with the ψ' function defined as

$$\psi'(x) = \psi(x) = \begin{cases} |x| & 0 \leq |x| < a \\ a & a \leq |x| < b \\ \frac{c - |x|}{c - a} & b \leq |x| < c \\ 0 & |x| \geq c \end{cases}$$

behaves universally well and also that some simple adaptive procedures such as choosing among trimmed means are much better than the simple ones, and usually better than the really adaptive procedures like Takeuchi's. Although there might be found still better estimators, the reviewer thinks that much overall improvement is impossible, and some simply adaptive procedures can be recommended to the practitioners. The interested reader will have to refer to the book about exactly what procedure he should choose.

The book contains many interesting details which should stimulate further investigations. And the reviewer would like only to add a few comments.

1) One should clearly distinguish the possible range of distributions, and the choice of "good" estimators depends on the size of the range. At least four cases should be distinguished: (i) from the normal to the t -distribution

with four to five degrees of freedom; (ii) to the t with two degrees of freedom and grossly contaminated normals; (iii) to the Cauchy; and (iv) to beyond the Cauchy. And for the first case the Hodges-Lehmann estimator seems to be better than this book suggests it to be. For the last case, which is rather unrealistic, we cannot do much better than simply use the sample median. For the third case some adaptive procedures should be used. The most difficult and practically most important case is the second one, and here the results of this book are most suggestive.

2) The desirable adaptive procedure should have the property that by applying it one not only can choose good estimates but can also get some idea how far the distribution is from the normal. In this sense test-estimation type procedures are better. Takeuchi's procedure can be modified in this way. In any case, a little loss in the efficiency can be disregarded if the "meaning" of the adaptive procedure is intuitively clear.

3) The location parameter problem is only a miniature of really relevant problems of statistical data analysis, and procedures which are directly applicable to a wider class of problems, for example to linear regressions, are more important than others. In this respect, M-estimators seem to be very promising.

K. TAKEUCHI

Faculty of Economics,
University of Tokyo, Tokyo, Japan

Insect Methodology

Aphid Technology. With Special Reference to the Study of Aphids in the Field. H. F. VAN EMDEN, Ed. Academic Press, New York, 1972. xiv, 344 pp., illus. \$18.50.

Many direct and indirect contributions to science and to society will result from studies undertaken in the International Biological Program. This book, the first on aphid technology, is one of these contributions.

Aphid Technology was conceived at a meeting chaired by Michael Way of Imperial College. Way and the other participants were intent upon establishing an I.B.P. undertaking on the biological control of aphids. It was obvious that a uniform technology would be necessary to operate such a project.

Under the editorship of H. F. van Emden, this book has gone far beyond

the original needs of the project. Twelve competent researchers have contributed to the book. Most of the chapters are so informative that even readers without access to the original publications can use the described techniques. Adequate bibliographies are included with each chapter. These are not as recent as might be desired. Of some 750 references, fewer than 30 are later than 1968 and half of these are in chapter 1. This, however, is not a serious defect because the authors selected well.

Information is presented that is not generally available because it has appeared in highly specialized publications. Fortunately some of the authors interpret, evaluate the methods, and add unpublished observations and experiences of their own and others. For example, I have used the technique for preserving aphids on slides that is suggested in chapter 1 and have experienced most of the difficulties listed. Here remedies are given for the difficulties.

It is estimated (table 1) that there are over 3700 species in this diverse group of insects. Despite the enormity of the task, the biological properties of the aphids are well covered and both the laboratory and the field methods used to obtain this information are given (chapter 2).

Aphids have natural enemies and, since outbreaks occur when these are interfered with or when populations temporarily outstrip them, methods for estimating the effectiveness of natural controls are included (chapter 4).

Many scientists are only beginning to realize the importance and the research possibilities of this group of insects. These researchers need to know how sampling is carried out (chapters 3 and 5), how the environment is measured (chapter 6), how problems of population dynamics are handled (chapter 7), and finally how the data obtained can be understood (chapter 8).

In this book specialists talk to each other, but it should be pointed out that if this were the only contribution the book made it would not be of wide enough interest for review in *Science*. The technology presented here can be applied to many invertebrate groups.

The insects are important. The methods are applicable to other animals. The book is recommended to aphid specialists and to other biologists.

JAMES B. KRING

Connecticut Agricultural Experiment
Station, New Haven