

lines change gradually in wavelength because the direction of gas flow, and hence the amount of Doppler shift, changes steadily. Margon pegged the period of this wobble in direction at 164 days.

Since most researchers focused their attention on the bizarre, moving spectral lines, they overlooked a big clue to the nature of the source that was hiding in the more normal emissions of the star. Even the apparently stationary spectral lines change in wavelength. David Crampton, Anne Cowley, and John Hutchings of the Dominion Astrophysical Observatory in Victoria, British Columbia, observed very slight Doppler shifts in these lines. The source approaches and then recedes at speeds up to 75 kilometers per second, says Crampton, and a full cycle takes about 13 days.

Such behavior is the signature of a binary, a two-star system similar to many of the strong x-ray sources in our galaxy. In this case, the two stars orbit each other in about 13 days.

But SS 433 does not behave like a normal x-ray binary. In addition to radiating some x-rays, although less than is typical in such binaries, it spends a lot of energy—roughly a million times the energy radiated by the sun—accelerating gas. The questions baffling theorists are why and how. At present, two very different mechanisms are considered promising.

With the discovery that SS 433 is a binary rather than a lone star, theorists breathed a sigh of relief. In the most popular type of model, which was first advocated by Martin Rees and Andy Fabian of Cambridge University, the high-speed gas streams away from SS 433 in two jets. Finding a source for the gas was a challenge that the star pair met easily. One of the stars sheds the gas, which its companion flings into space. The companion is believed to be either a neutron star or a black hole.

In many jet models, the reason gas is ejected from the system is that too much is falling toward the compact companion. Normally, material flowing between

the two stars would be converted into x-rays and added to the superdense star. But in the case of SS 433, so much mass is transferred that x-rays are extremely intense. This radiation “pushes the infalling matter up faster than gravity pulls it down,” explains Jonathan Katz of UCLA. The only place the expelled gas can go is out of the plane containing the two stars and all the material flowing between them. It travels just fast enough (about one-quarter the speed of light) to escape the gravitational pull of the compact object. According to Craig Sarazin of the University of Virginia, about 1000 times as much mass is transferred in SS 433 than is normal in x-ray binaries.

Theorists have offered several possible explanations for the beams’ wobbling. Strong tidal interactions between the two stars could cause the wobble, say Edward van den Heuvel of the University of Amsterdam, Jeremiah Ostriker of Princeton University, and Jacobus Pettersen of the University of Illinois, Urbana. Alternatively, Mitchell Begel-

Carbon Budget Not So out of Whack

Every year plants exchange more than 100 billion metric tons of carbon with the atmosphere in the form of carbon dioxide. The ocean spews forth and reabsorbs almost as much carbon dioxide as it churns up its deepest waters and as its surface waters warm and cool with the seasons. Even these mammoth quantities seem insignificant next to the thousands of billions of tons of carbon stored in the oceans, plants, and soil of Earth.

The task of measuring small changes in these flows of carbon dioxide through the atmosphere is difficult but necessary. Predictions of the possibly catastrophic climatic warming that could result from man’s increasing the carbon dioxide content of the atmosphere require such measurements. Among those trillions of tons of carbon, researchers are trying to determine whether human activities each year are releasing only 5 billion tons (5 gigatons) or perhaps as much as 10 or 15 gigatons of carbon as carbon dioxide to the atmosphere. Everyone agrees that an amount equivalent to about half of the 5 gigatons of fossil fuel carbon now burned each year ends up in the atmosphere as carbon dioxide. Many oceanographers have thought that most of the rest dissolves in the surface waters of the ocean. That would leave all of the anthropogenic carbon dioxide accounted for and the cycle balanced, except that some terrestrial biologists suggested a few years ago that the destruction of forests, especially the increasing exploitation of the tropical forests, adds even more carbon dioxide to the cycle (*Science*, 30 September 1977, p. 1352). Initial estimates of the land’s contribution ranged as high as twice the size of the fossil fuel source, but oceanographers recoiled at the suggestion of putting all of that carbon into the ocean. It could not be done, they said.

Few new field data have been gathered, and the oceanog-

raphers and biologists still do not agree, but more leisurely analyses of the available data have narrowed the range of disagreement, primarily by shrinking the possible terrestrial source. In one recent study of the biosphere’s role in the carbon cycle, Wolfgang Seiler of the Max Planck Institute for Chemistry in Mainz and Paul Crutzen of the National Center for Atmospheric Research in Boulder concluded that the land may be acting as either a small source of carbon dioxide or as a small sink by absorbing more carbon dioxide than it gives off.

In particular, Seiler and Crutzen considered how fires, such as those used in the slash-and-burn agriculture of the tropics, wild fires, the burning of wood for fuel, and the clearing of land for urbanization, could affect the conversion of plants into carbon dioxide. For the first time, Seiler and Crutzen tried to account for the carbon preserved as charcoal following a fire. They estimated that this decay-resistant form of carbon might provide a sink, rather than a source, for 0.5 to 1.7 gigatons of the carbon burned each year. Taking account of other losses and gains, such as the oxidation of organic soils exposed by farming and reforestation of cleared areas, they concluded that the land could be either gaining or losing as much as 2 gigatons of carbon each year.

Although it is a significant fraction of the 5 gigatons released from fossil fuels, a maximum of 2 gigatons per year is small compared with the estimate of 4 to 8 gigatons per year (with a possible range of 2 to 18 gigatons) offered several years ago by George Woodwell of the Marine Biological Laboratory and his colleagues. But that figure too may be shrinking. In a summary paper presented at a Department of Energy research conference in Washington in May, it was reported that Woodwell’s current best, though

mann of the University of California, Berkeley, and Sarazin speculate that the wobble may be due to relativistic effects that warp space near the compact object.

The jets from SS 433 may be narrow because the gas squirts out through "nozzles," say Davidson and Richard McCray of the Joint Institute for Laboratory Astrophysics in Boulder. According to their model, excess mass shed by the more normal star spirals toward, but does not reach, the compact object. ("I believe it is a neutron star, but Kris seems to favor a black hole," says McCray. Even collaborators do not agree completely on SS 433.) Instead the gas forms a dense cloud, the inner edge of which is shaped like two nozzles aimed in opposite directions out of the plane of the two stars. "Gas evaporates into the nozzles," says Davidson, where it is accelerated outward by the pressure of radiation.

In the nozzles the streaming gas is far too hot for any neutral hydrogen to exist and emit the spectral lines that won SS

433 its fame. After emerging from the nozzles, however, the gas cools, and hydrogen ions combine with electrons.

"The reason the jets emit strongly is that they are rubbing against a stellar wind from the normal star," explains McCray. Each atom is ionized repeatedly by friction between the jets and the much slower gas of the wind. Each time an ion recombines with an electron, the newly formed atom emits the characteristic radiation as the electron cascades toward its ground state. McCray estimates that the region where this process occurs is about 1 billion kilometers long.

Both the means of accelerating the gas and its direction of motion set one model apart from the popular jet models. George Collins II, Gerald Newsom, and Richard Boyd of Ohio State University propose that the strong magnetic field of one of the stars, rather than the pressure of radiation, pushes the emitting gas. And the gas, riding the rotating magnetic field, whizzes around the binary, not away from it.

According to the Ohio State model, streams of ions blowing out from the magnetic poles of one of the stars collide in two regions on opposite sides of the binary. The moving spectral lines are emitted by the high-density, cool gas where the streams merge. As the star spins, the emitting regions circle the binary to trace out a ring. Since the ring is huge—30 times the diameter of the solar system, says Collins—and the gas takes about 164 days to orbit the binary, it cruises faster than one-quarter the speed of light.

The sheer size of the ring has implications that may allow this model to be tested by observations. Light emitted from the far side of the ring takes about 4½ days longer to reach the earth than light emitted from the near side. Thus signals emitted simultaneously from the two emitting regions may be received at different times by telescopes. The time lag will vary as the two regions circle around the ring. According to Collins, a preliminary analysis of published data

not final, estimate for the 1970 release is 2 to 4 gigatons of carbon. "We can't rule out the 8 gigatons per year yet," says Woodwell, "but I would guess, and it's only a guess, that it's less than 8 and more than 2 gigatons per year. We're certainly ruling out the extreme number of 18."

Many terrestrial biologists are thinking about smaller releases of about 2 gigatons per year, citing a number of recent studies that lend support for a smaller contribution from the land. For example, Jerry Olson of the Environmental Sciences Division of the Oak Ridge National Laboratory estimates that living land plants contain only 560 gigatons of carbon rather than the once generally accepted 800 gigatons, which would mean lower fluxes of carbon dioxide from the exploitation of forests. The mass of plants actually being converted into carbon dioxide might be even smaller, according to Charles Hall of Cornell University, because as much as half of the forest being cleared in the tropics may not be dense virgin stands of trees but rather thinner stands that grew back since an earlier clearing. In addition, the carbon dioxide that is being released in the tropics is probably partially compensated for by the growth of temperate forests, according to the final report of a workshop conducted by the Institute of Ecology in Indianapolis. The temperate forests, which were extensively cleared in the 19th century, now appear to be regrowing fast enough to store about 1 gigaton of carbon per year, according to the workshop report.

While the biospheric source has been shrinking, oceanographers have continued to refine their estimates of how fast the ocean can absorb carbon dioxide from the atmosphere. Although oceanographers no longer suggest that the terrestrial biosphere must absorb much carbon dioxide, they cannot find much room in the ocean to place carbon

dioxide from a terrestrial source. By tracing the path of nuclear bomb debris from the surface into deeper waters, and by using other techniques, Wallace Broecker of the Lamont-Doherty Geological Observatory has calculated that about 37 percent of the fossil fuel carbon dioxide released since 1958 could have been absorbed. Although Broecker and his colleagues are still adamant about the impossibility of dissolving several extra gigatons of biospheric carbon in the ocean, Broecker concedes that as much as 0.5 gigaton per year from a terrestrial source might be accommodated in the ocean. "One gigaton would be pushing it, but I suppose it could be possible," he says. Robert Bacastow of Scripps Institution of Oceanography reports that the ocean models he and Charles Keeling of Scripps work with do not allow a much larger terrestrial source than Broecker's does.

Although oceanographers and terrestrial biologists are beginning to talk about numbers in the same general range, a gap remains and errors in the estimates are still large. Some possible small sinks for carbon dioxide that could help close the gap are now being given serious consideration. The oceanographers will also be collecting crucial new data next year in the northern North Atlantic under the Transient Tracers program. No equivalent program for gathering data from the more complex terrestrial biosphere exists yet. The counteracting effects over the last few decades of resurging temperate forests and accelerating tropical forest clearing may have minimized the role of the terrestrial biosphere in the carbon dioxide problem, but researchers now realize that a land source could become a major factor as increasing population and the increasing cost of fossil fuels put greater pressure on the world's forests.—RICHARD A. KERR