survival, a finding which came as a surprise at a time when transfusion was considered a contraindication for kidney graft candidates.

Are we not forgetting the pioneers?

MARILENA FOTINO

New York Blood Center, 310 East 67 Street, and Cornell University Medical College, New York 10021

References

 G. Opelz, D. P. S. Sengar, M. R. Mickey, P. I. Terasaki, *Transplant. Proc.* 5, 253 (1973); G. Opelz and P. I. Terasaki, *Lancet* 1974-II, 696 (1974).

With respect to Marx's article of 8 August, it should be noted that there is still isolated disagreement as to the beneficial effect of transfusion before transplant on survival of kidney grafts from cadaver donors. An analysis (1) of the transfusion effect on 514 transplants performed at seven centers in Illinois demonstrates that recipients who were given transfusions before transplant experienced fewer early graft failures than those who were not. However, the Illinois study also demonstrated an increase in late graft failure in recipients who were given such transfusions, leading to a 2- to 7year cumulative graft survival that was not different in recipients who received transfusions before transplant and those who did not.

In our own small transplant series from cadaver donors (82), we have not been able to discern any beneficial effect of transfusion before transplant after 1 or 4 years of follow-up (among recipients who were given such transfusions, graft survival after 1 year was 75 percent and after 4 years, 53 percent; among recipients who were not, it was 70 percent and 50 percent, respectively). We certainly do not find the data in favor of transfusion before transplant "overwhelming." We electively give transfusions to all patients during transplant operations but find no justification for deliberately giving transfusions before a transplant.

P. R. McConnachie

Memorial Medical Center, Springfield, Illinois 62702

A. G. BIRTCH

Southern Illinois University School of Medicine, Springfield 62708

References

1. O. M. Jonasson and M. F. Mozes, personal communication.


Erratum: In the report by R. Ganapathy (22 Aug., p. 921), Pd should have been 2 instead of 20 on line 2 in the middle column of p. 922, and Au should have been 3×10^{-2} instead of 3×10^{-1} on line 6 in the same column.

Erratum: In the article by R. Hekinian et al. (28 Mar., p. 1433), in line 3 of the caption for Fig. 7 (p. 1440), "Fe-rich clay-like material" should have read "Fe-bearing opaline material."



Circle No. 108 on Readers' Service Card

How To Gain Control Of Your Environmental Control Problems:



Call Sherer.

Our engineers and designers can help you select the optimum environmental chamber for your particular needs in the control of light, temperature and humidity.

We offer a complete line of table-top, reach-in and walk-in chambers for incubation and plant, animal and tissue studies.

 Due to their unique design and construction, they can be quickly

installed and easily adjusted to your individual needs. And to accommodate the changing requirements of scientific inquiry, Sherer can also provide you with tailor made environmental "rooms," in 96 standard models and virtually hundreds of modifications.

So call or write today for your copy of our new catalog. And find out how to gain control of your environmental control problems.

Rheem Manufacturing Company, Refrigeration Products Division, 1100 Memorial Drive, West Columbia, South Carolina 29169, Telephone 803-796-1700, TWX 810-666-2103.

SHERER

Circle No. 150 on Readers' Service Card