bioluminescent and nonbioluminescent Lampyridae, active at dusk. However, the present combination of electrophysiological, spectroscopic, and field behavioral data strongly support the hypothesis that the shift in color of firefly species bioluminescence and in visual spectral sensitivity from green to longer wavelengths is an evolutionary adaptation to increase the efficiency of sexual signaling in different, prevailing light environments.

> ABNER B. LALL HOWARD H. SELIGER WILLIAM H. BIGGLEY

McCollum-Pratt Institute and Department of Biology, Johns Hopkins University, Baltimore, Maryland 21218 JAMES E. LLOYD

Department of Entomology, University of Florida, Gainesville 32611

References and Notes

- 1. J. E. Lloyd, in *Bioluminescence in Action*, P. J. Herring, Ed. (Academic Press, New York, 1978), p. 570 and references therein. York,
- Sexual responses can be elicited in the field from both sexes with various light sources (flashlights, with or without colored filters; cigarettes; matches; glowing light organs of other species) provided the stimulated signal has the correct timing and there is some portion of the spectrum of the artificial source that overlaps the effective visual spectral sensitivity. The photon intensity for the bioluminescent
- flash is low, and for dark-adapted vision is below the threshold of human cones for precise color discrimination. Therefore spectrophotometric measurements in the field are essential for determeasurements in the held are essential for determining the color of the bioluminescent signal [W. H. Biggley, J. E. Lloyd, H. H. Seliger, J. Gen. Physiol. 50, 1691, 1967; W. D. McElroy, H. H. Seliger, M. DeLuca, in The Physiology of Insecta, M. Rockstein, Ed. (Academic Press, New York, 1974), vol. 2, p. 411].
 4. H. H. Seliger, Photochem. Photobiol. 21, 355 (1975)
- (1975). W. D. McElroy and H. H. Seliger, in Molecular 5. W. D. McErroy and H. H. Scinger, in Molecular Architecutre in Cell Physiology, T. Hayashi and A. Szent-Gyorgyi, Eds. (Prentice-Hall, Engle-wood Cliffs, N.J., 1966), p. 63. J. E. Lloyd, in How Animals Communicate, T.
- Sebeok (Indiana University Press, Blooming-ton, 1977), p. 1128. Dartnall (1953) proposed that the absorption bandwidth of all visual pigments when plotted
- on a frequency scale were identical, and inde-pendent of the absorption maxima of the pigment; he devised a constant bandwidth template now called the Dartnall nomogram. However, systematic deviations from this nomogram, with short wavelength pigments having broader short wavelength pigments having broader bandwidths and longer wavelength pigments, narrower bandwidths, made Ebrey and Honig (1977) propose three different templates for pig-ments absorbing in the short, middle, and longer wavelengths [H. J. A. Dartnall, Br. Med. Bull. 9, 24 (1953); T. G. Ebrey and B. Honig, Vision Res. 17, 147 (1977)]. Sameitivity is the reciprocal of the quanta needed
- 8. Sensitivity is the reciprocal of the quanta needed Scisibility is the recipical of the quanta needed to evoke a criterion response. The details of the method, as well as the experimental set up are given in A. B. Lall, R. M. Chapman, C. O. Trouth, J. A. Holloway, J. Comp. Physiol. 135, 21 (1980).
- 9. The standard deviation of an experimental determination of the peak wavelength of emission sion spectrum from a live firefly is approximately 1.2 nm(3), while that for the ERG spectral sensi-tivity determination in this work is 3 nm. Since, in the measurements reported here, the number of samples for each of the measurements ranged from N = 2 to N = 12 the 95 percent confidence limits for the specification of identity between an ERG peak and a bioluminescence emission peak re less than 2.3 nm.
- are less than 2.3 nm. 10. All insects possess sensitivity maximums in the green ($\lambda_{max} = 480$ to 550 nm) [T. H. Goldsmith and G. D. Bernard, in *The Physiology of Insecta* R. M. Rockstein (Academic Press, London, 1974), vol. 2, p. 165]. Some butterflies [G. D.

Bernard, Science 203, 1125 (1979)] and the moth Spodoptera example 105 (17) and the month Spodoptera example (H. Langer, B. Hamann, C. C. Meinecke, J. Comp. Physiol. 129, 235 (1979)] have additional red receptors.

 The late flashing *Photuris aureolucens* (554 nm) in the same habitat initiates flashing activity 20 minutes earlier than its sibling species Photuris caerulucens (551 nm), while among the early flashers Photinus macdermotti (Maryland) (570 nm) initiates flashing activity earlier than P. macdermotti (Florida) (565 nm); P. scintillans (578 nm) in the same habitat initiates flashing activity 20 to 25 minutes earlier than P. pyralis

(563 nm). There are no known reverse cases. Research supported in part by Department of Energy contract EVO-3277 (to H.H.S.), NSF grant DEB 792-1744 (to J.E.L.), and grants from 12. the Kettering Family Foundation and Robert L. Conway (to A.B.L.). Contribution number 1054 of the McCollum-Pratt Institute and the Department of Biology, Johns Hopkins University, Baltimore, Md

23 November 1979; revised 28 April 1980

Induction of Suspension Feeding in Spionid Polychaetes by **High Particulate Fluxes**

Abstract. The feeding behavior of three species of spionid polychaetes varied with water velocity. At moderate flows the worms ceased deposit feeding, formed their feeding tentacles into helices, and lifted them into the water column to capture material in suspension. This behavior was apparently a response to increased flux of suspended matter at high flows rather than to flow velocity alone. Organisms capable of switching their feeding behavior may be common in dynamically variable benthic environments.

Deposit-feeding polychaete worms are a major group in soft-bottom areas of both the shallow and deep sea (1). These sediment-inhabiting annelids ingest food materials found on the sediment surface or within the sediment. Deposit feeders in general are considered to exploit a different food source than benthic suspension feeders, which obtain their food by removing it from the overlying water. Indeed, differences in factors controlling food supply are one explanation for the alternative development of predominantly deposit- or suspension-feeding benthic communities (2). Higher water velocities increase the flux of suspended materials, favoring suspension feeders, while removing fine-grained particulate matter from the sediment or preventing the deposition of this material, to the disfavor of deposit feeders.

A clear-cut distinction between these two feeding modes is probably unrealistic. Because there can be considerable resuspension of bottom materials by water currents in both the shallow (3) and deep sea (4), it could be advantageous for an animal to adjust its feeding behavior to utilize food in suspension or deposited on the bottom, depending on the



Fig. 1. Tentacles of Pseudopolydora kempi japonica, held in characteristic helical arrangement during suspension feeding. The current is moving from left to right at about 13 cm/sec (×4).

562

dominant source of food at a particular time.

Spionid polychaetes are a widely distributed benthic group (5). They are usually considered to be deposit feeders, although facultative suspension feeding is known for this family (6, 7). We report a novel suspension feeding mechanism in three species of spionid polychaetes. Apparently the mechanism is employed in response to the increased flux of suspended materials occurring at high water velocities.

A seawater flume at the Friday Harbor Laboratories provided controlled flow velocities (8). The flume operated with continuously supplied Puget Sound seawater or with a pump and filter system that recirculated the water while removing particulate matter (9). A recessed box in the flume bottom contained the sediment and animals; the sediment-water interface was coplanar with the flume bottom. Pseudopolydora kempi japonica, Boccardia proboscidea, and Pygospio elegans were collected from the intertidal zone of False Bay on San Juan Island, Washington, by sieving sediments through a screen with 500- μ m mesh. The sieved sediment, free of macrofauna, was placed in a flume box. Worms, sorted and placed on the sediment surface, usually burrowed and established new tubes within 1 hour. Experiments were performed with one species at a time. Water velocity was measured with a commercial quartzcoated hot film (TSI, Inc., model 1231W) operated at 20°C overheat by a TSI anemometer (8). Vertical position of the probe was determined by a precision vernier. Mean velocities were calculated



Fig. 2. Proportion of *P. kempi japonica* deposit feeding (\Box) or displaying helical tentacles (\bullet) versus water velocity 0.4 cm above the bed. Each pair of data points represents an average for 90 worms. Numbers in the body of the figure are numbers of replicate observations falling at the same point.

from 30-second analog averages of the d-c voltages. Concentrations of suspended particulates were measured by filtering 1.0-liter samples of seawater (collected at the outflow flume end) through tared Millipore HAWP filters (nominal pore size, 0.45 μ m), rinsing with filtered distilled water to remove salts, drying at 60°C, and weighing to the nearest 0.01 mg.

Spionid polychaetes possess two anterior feeding tentacles. During feeding, the tentacles are extended from the tube and, during conditions of low flow (0 to 2 cm/sec) (10), held on or near the sediment surface. Food particles are picked up near the distal ends of the tentacles and transported to the mouth via a ciliated groove. At higher water velocities (~ 5 cm/sec), the worms lift their tentacles into the water column and extend them downstream, with the axes parallel to each other and to the bed. The tentacles are twisted into gradually tapering helices (Fig. 1).

Behavior was observed under low magnification while the water flow through the flume was varied. Results for *P. kempi japonica* are given in Fig. 2. (Results for *B. proboscidea* and *P. elegans* are qualitatively similar; however, the larger size and greater abundance of *P. kempi japonica* make it best suited for quantitative observations.) As water velocity increased, the proportion of *P. kempi japonica* feeding on deposited particles decreased (Spearman's $\rho = -.50$, P < .05), while an increasing proportion extended their tentacles in the helical arrangement ($\rho = .39, .10 < P < .15$) (*11*).

Since the helical shape increases the surface area of the tentacle normal to the flow, it seems likely that the worms were suspension feeding. To see whether the worms were capable of this behavior, we introduced newly hatched brine shrimp upstream of the sediment box. The shrimp were captured along the entire length of the tentacles, transported to the mouth by ciliary action, and ingested. Natural flocculent material was also captured. Dimensionless indices for capture modes by a single fiber (as a spionid tentacle may be considered) can be computed according to formulas discussed by Rubenstein and Koehl (12). The values for the brine shrimp experiment are (i) effectiveness of capture due to direct interception, approximately 1.0; (ii) effectiveness of capture due to inertial impaction, approximately 0.009; and (iii) effectiveness of capture due to gravitational deposition, approximately 0.003. These calculations and the visual observations suggest that direct interception by the tentacle is the primary means by

which these annelids feed on larger particles in suspension. However, the importance of inertial impaction may be underestimated if the helical tentacles impart centrifugal forces to the water passing through them.

To determine whether the helical form of the tentacles was controlled by the worms or formed passively during high flow due to some structural characteristic, several animals were narcotized in 7.5 percent MgCl₄ and implanted in the sediment with their tentacles exposed to the flow. At water velocities that induced control worms to hold their tentacles helically, the narcotized specimens' tentacles trailed limply downstream. A great deal of control over tentacle shape and positioning is undoubtedly exerted by the worm. For example, as water velocity increased, the number of complete turns made by the tentacles also increased (13), adding to the effective filtering surface at higher particulate fluxes. This level of control may not be typical of other deposit-feeding polychaetes, such as ampharetids and terebellids, which do not show comparable tentacle musculature (14) and which have difficulty feeding at comparable flow velocities (15).

To determine whether the helical arrangement was a response to the increased flux of suspended material that occurs at high water velocities or a re-



Fig. 3. Feeding behavior as a function of particulate matter flux. Closed bars represent particulate flux; open bars, number of worms displaying helically arranged tentacles; crosshatched bars, number of worms deposit feeding. The number of worms feeding with extended tentacles was determined every 30 minutes. Particulate flux was calculated by multiplying the water velocity (12 cm/sec) by the suspended particulate matter concentration (determined from the median of three 1.0liter water samples taken for each calculation). The water samples were obtained at time 0, and then the pump and filter system was engaged. Two hours later, water samples were taken before the pump and filter were disengaged, and the flume was again operated with continuously renewed water from the laboratory seawater system. Samples of this water were taken 1 hour later. Worms responded gradually to engagement or disengagement of the filter because of the large volume of the flume system (~ 270 liters).

sponse to higher water velocity per se, the flume's pump and filter system were used to vary the concentration of suspended particulate matter (Fig. 3). By the time approximately 40 percent of the particulates (by weight) was filtered out, most worms had withdrawn their tentacles into their tubes. When the experiment was repeated with the filter element removed from the recirculation system, the concentration of suspended particulate matter (and hence the particulate flux) remained constant, as did the number of worms actively feeding. Thus it was not the operation of the pump and filter that inhibited feeding behavior, but the reduced particulate flux.

It has been known for some time that certain benthic organisms, including some spionids, nereids, fabricine sabellids, and oweniids among the polychaetes (6), tellinids among the bivalves (16), and amphipods among the crustaceans (17) are capable of both suspension and deposit feeding. The growing understanding of benthic boundary layer dynamics and rapidly improving flow measurement technology now permit an analysis of the factors influencing the preferred feeding mode. We suggest that animals capable of switching their feeding behavior typify environments of rapidly varying flow characteristics. Rather than being a troublesome anomaly in trophic classifications (6), such animals form a distinctive indicator group of their own.

GARY L. TAGHON ARTHUR R. M. NOWELL PETER A. JUMARS Department of Oceanography,

University of Washington, Seattle 98195

References and Notes

- R. R. Hessler and P. A. Jumars, *Deep-Sea Res.* 21, 185 (1974); P. A. Jumars and K. Fauchald, in *Ecology of Marine Benthos*, B. C. Coull, Ed. (Univ. of South Carolina Press, Columbia, 1077)

- (Univ. of South Carolina Press, Columbia, 1977), p. 1.
 2. H. L. Sanders, Bull. Bingham Oceanogr. Collect. 15, 345 (1956); D. J. Wildish, Helgol. Wiss. Meeresunters. 30, 445 (1977).
 3. M. R. Roman and K. R. Tenore, Estuarine Coastal Mar. Sci. 6, 37 (1978); D. K. Young, Vie Milieu Ser. A 22 (Suppl.), 557 (1971).
 4. E. T. Baker and R. A. Feely, Science 200, 533 (1978); P. E. Biscaye and S. L. Eittreim, Mar. Geol. 23, 155 (1977); R. A. Kerr, Science 208, 484 (1980).
 5. K. Fauchald. The Polychaete Worms (Natural)
- . Fauchald, The Polychaete Worms (Natural History Museum of Los Angeles County, Los
- Angeles, 1977), p. 22.
 and P. A. Jumars, Oceanogr. Mar. Biol. Annu. Rev. 17, 193 (1979).
 D. M. Dauer, C. A. Maybury, R. M. Ewing, un-
- published manuscript. A. R. M. Nowell, P. A. Jumars, R. F. L. Self in 8.
- preparation. The filter removes particles > 10 μ m in size. All water velocities are for 0.4 cm above the bed, the approximate height at which the helical 10 tentacles are held.
- Proportions at any given water velocity do not always sum to 1.0 because a few worms usually engaged in other activities (such as tube build-
- ing). D. I. Rubenstein and M. A. R. Koehl, Am. Nat. 12. 111, 981 (1977).

- 13. When the water velocity is 5 cm/sec, the median number of turns in a helical tentacle is 1 (N = 10); at 12 cm/sec, 1.25 (N = 8); at 19 cm/ ec, 2.25 (N = 6); and at 26 cm/sec, The trend of increased coiling of the tentacles as velocity increases is highly significant (P << .01, Jonckheere trend test) [A. R. Jonckheere, Biometrika 41, 133 (1954)]. 14. R. P. Dales, J. Mar. Biol. Assoc. U.K. 34, 55
- 1955)
- 15. In comparable experiments with Hobsonia florida (Ampharetidae) and Thelepus crispus (Tere-bellidae), increased flow velocities simply reduced the proportion of individuals successfully feeding.
- 16. A. E. Brafield and G. E. Newell, J. Mar. Biol. Assoc. U.K. 41, 81 (1961); R. N. Hughes, *ibid.* 49, 805 (1969).
- 17. L. Mills, J. Fish. Res. Board Can. 24, 305 (1967).
- 18. We thank the director of the Friday Harbor Laboratories, University of Washington, for providing laboratory space and equipment. We also thank M. A. R. Koehl and S. Vogel for helpful Suggestions and J. A. Taghon for photography. Supported by DOE contract DE-AT06-76-EV-75026 (DE-EV-75026-86) and ONR grant N00014-75-C-0502.

19 May 1980

Hormone Accumulation in a Sexually Dimorphic Motor Nucleus of the Rat Spinal Cord

Abstract. The fifth and sixth lumbar segments of the rat spinal cord were found to contain a sexually dimorphic nucleus, the spinal nucleus of the bulbocavernosus (SNB). The SNB, which contains motoneurons innervating perineal striated muscles in normal male rats, is diminished or absent in normal females and in males with a genetic mutation rendering them insensitive to androgens. The presence of the nucleus is apparently not dependent on genetic sex, but on the action of androgens. The motoneurons of the adult male SNB accumulate hormone after systemic injections of radioactive testosterone or dihvdrotestosterone, but not estradiol, and the SNB motoneurons accumulate more of the injected androgens than do other motoneurons in the same spinal segments. These results demonstrate a morphological sex difference in hormone-sensitive motoneurons that are probably involved in the sexually dimorphic copulatory behavior of the rat.

Because of the obvious sex differences in the behavior of vertebrates, one might expect to find differences in the central nervous system (CNS) of the two sexes. Raisman and Field (1) reported that in rats the "strial portion" of the preoptic area (POA) has more synapses from nonamygdaloid sources in females than in males. Other anatomical sex differences described in the nervous system include the number of autonomic preganglionic neurons in cats, and the dendritic field patterns in the POA of hamsters (2). Nottebohm and Arnold (3) reported gross sexual dimorphism in the size of certain brain nuclei of song birds. These nuclei are larger in males and are known to play a role in the singing behavior which only males display. Gorski et al. (4) found a strikingly dimorphic nucleus in the rat POA which is larger in males than in females. Because the last two dimorphisms are easily detected, they provide convenient measures of the process of sexual differentiation of the brain, and therefore study of these systems may lead to a better understanding of the factors critical to sexual differentiation of vertebrate behavior. We now report a prominent dimorphism in a motor nucleus of the rat spinal cord, the neurons of which accumulate radioactivity after injections of tritiated androgens but not estradiol. This dimorphism is useful for studying sexual differentiation of the CNS because it offers advantages dis-

0036-8075/80/1031-0564\$00.50/0 Copyright © 1980 AAAS

tinctive to motoneurons, for example, readily determined behavioral function, electrophysiological accessibility due to the large somas, relatively simple inputs and outputs, and ease of study in early development.

Motoneurons were identified by injecting horseradish peroxidase (HRP) into each of the three striated muscles attached to the rat penis: the ischiocavernosus (IC), bulbocavernosus (BC), and levator ani (LA) (5). The muscles of 30 male anesthetized Sprague-Dawley rats were injected with 3 to 50 μ l of a 30 percent solution of HRP (Sigma type VI) in saline by means of a $10-\mu$ l syringe under a dissecting microscope (6). Multiple injections were used to distribute the HRP evenly throughout the muscle, and afterward the exposed perineal region was flushed with saline (7). The animals were killed 24 hours later and the retrogradely transported HRP was stained in the spinal cord segments caudal to thoracic 13 (8).

After the injection of HRP into IC, retrogradely labeled motoneurons were found in the extreme ventrolateral quadrant of the ventral horn in the fifth and sixth lumbar spinal segments (L₅ and L_6). Injection of HRP into either the BC or LA resulted in labeled cells being found in the dorsomedial portion of the ventral horn, 50 to 250 μ m from the midline and 200 to 400 μ m below the ventral margin of the central canal (Fig. 1a, ar-