

SCIENCE

Published by the **American Association for the Advancement of Science (AAAS)**, *Science* serves its readers as a forum for the presentation and discussion of important issues related to the advancement of science, including the presentation of minority or conflicting points of view, rather than by publishing only material on which a consensus has been reached. Accordingly, all articles published in *Science*—including editorials, news and comment, and book reviews—are signed and reflect the individual views of the authors and not official points of view adopted by the AAAS or the institutions with which the authors are affiliated.

Membership/Circulation

Director: Michael Spinella

Fulfillment: Marlene Zendell, *Manager*; Gwen Huddle, *Assistant Manager*; Mary Curry, *Member Service Supervisor*; Pat Butler, Helen Williams, Robert Smariga, *Member Service Representatives*

Promotions: Dee Valencia, *Manager*; Laurie Baker, Hilary Baar, *Assistants*

Research Manager: Kathleen Markey

Financial Analyst: Jacquelyn Roberts

Administrative Assistant: Nina Araujo de Kobes

Advertising and Finance

Associate Publisher: Beth Rosner

Advertising Sales Manager: Susan A. Meredith

Display Recruitment Sales Manager: Janis Crowley

Financial: Deborah Rivera-Wienhold, *Manager*; Julie Eastland, *Senior Analyst*; Andrew Joyce, *Junior Analyst*

Marketing Manager: Laurie Hallowell

Traffic Manager: Tina Turano

Traffic Manager (Display Recruitment): Daniel Moran

Line Recruitment: Michele Pearl, *Manager*; Millie Muñoz-Cumming, *Assistant*

Reprints Manager: Corrine Harris

Permissions Manager: Arlene Ennis

Advertising Assistants: Allison Pritchard, Kelly Nickerson, Debbie Cummings

Send materials to *Science* Advertising, 1333 H Street, NW, Washington, DC 20005, or FAX 202-682-0816.

SALES: Northeast/E. Canada: Fred Dieffenbach, Rt. 30, Dorset, VT 05251; 802-867-5581, FAX 802-867-4464 • **Mid-Atlantic:** Richard Teeling, 28 Kimberly Place, Wayne, NJ 07470; 201-904-9774, FAX 201-904-9701 • **Southeast:** Mark Anderson, 1915 Brickell Avenue, Ste. CC-1, Miami, FL 33129; 305-856-8567, FAX 305-856-1056 • **Midwest:** Donald Holbrook, 1110 North Harvey, Oak Park, IL 60302; 708-386-6921, FAX 708-386-6950 • **West Coast/W. Canada:** Neil Boylan, 2847 Fillmore, Ste. 3, San Francisco, CA 94123; 415-673-9265, FAX 415-673-9267 • **Germany/Switzerland/Austria:** Ric Bessford, World Media Services, Leopoldstrasse 52, 8000 Munich 40, Germany; +49-089-39-00-55, FAX +49-089-39-00-15 • **Japan and Far East:** Masy Yoshikawa, Orient Echo, Inc., 1101 Grand Maison Shimomiyabi-cho 2-18, Shinjuku-ku Toyko 162, Japan; +3 3235-5961, FAX +3 3235-5852 • **UK, Scandinavia, France, Italy, Belgium, The Netherlands:** Andrew Davies, 1 Newbridge View, Micklehurst Road, Mossley, Ashton-under-Lyne, OL5 9SE, Great Britain; +44-457-838-519, FAX +44-457-838-898 • **Other:** Contact *Science* Advertising: 202-326-6544, FAX 202-682-0816.

Information to Contributors appears on pages 36–38 of the 3 January 1992 issue. Editorial correspondence, including requests for permission to reprint and reprint orders, should be sent to 1333 H Street, NW, Washington, DC 20005. **Science Telephone:** 202-326-6500. London office: 071-435-4291. **Subscription/Member Benefits Questions:** 202-326-6417. **Other AAAS Programs:** 202-326-6400.

LETTERS

Tree-Ring Research

Regarding a photograph by one of us (B.M.B.) that appeared with the article "There's a new offering on campus: Global change 101" by Ann Gibbons (*Research News*, 22 May, p. 1146), I would like to clarify a few points. First, the photograph shows Mike Peterson of the Tasmania Forestry Commission and Mike Barbetti of the University of Sydney working near the summit of Mount Read, in a group of trees killed by fire in 1961. Neither Peterson nor Barbetti is a Lamont-Doherty researcher, as stated in the caption; both were assisting on an ongoing project directed by Ed Cook, a research scientist at Lamont-Doherty. We cut and sectioned this standing, dead tree for our study; but the rest of our samples were taken with the use of increment corers, which remove only a 4-millimeter dowel of wood from the tree and cause no discernible harm to living trees. We do not usually cut trees, and when we do only dead trees are cut. Additional samples taken at this site included buried and partially buried, nonliving stems and eroded stumps from within the fire-killed site only, and not from the adjacent, living site. We make every effort to have minimal impact in the areas in which we work, while at the same time getting the best samples for our research interests.

Brendan M. Buckley
Edward R. Cook
Gordon C. Jacoby

*Tree-Ring Laboratory,
Lamont-Doherty Geological Observatory
of Columbia University, Palisades, NY 10964*

Evolution of HIV in Africa

Steve Sternberg's *Research News* article of 15 May (p. 966) describes a phylogenetic analysis which concludes that West Central Africa may contain a lineage of HIV-1 (human immunodeficiency virus-1) that is ancestral to the other HIV-1 isolates analyzed to date. This conclusion has been criticized on the basis of the relatively low prevalence of HIV-1 in that region of Africa, but this low prevalence is entirely consistent with the patterns expected from current evolutionary considerations of HIV virulence (1). These evolutionary considerations suggest that the pandemic HIV evolved to a high level of virulence in

mid-Central and East Africa in response to an increase in multiple-partner sexual contact brought about by the socioeconomic crisis that occurred there during the 1970s. The lower rates of sexual contact along the west coast of Africa should have favored the evolution of lower virulence. Accordingly, current evidence indicates that the HIV-2 in West Africa has a lower virulence and lower prevalences than the pandemic HIV-1 (1–3). The limited comparisons to date also suggest that these low prevalences of HIV-2 are remaining fairly stable in areas of West Africa where HIV-2 predominates (4). According to this evolutionary argument, the same should be true of HIV-1 strains that have evolved endemically in areas with relatively low rates of multiple-partner sexual contact. That is, we can expect such strains of HIV-1 in western regions to have a lower, more stable prevalence than the HIV-1 strains in mid-Central and East Africa.

This evolutionary argument can be tested by assessing the virulence of the root West African HIV-1 lineages in humans by quantifying, for example, the time between the onset of infection and the onset of AIDS. The evolutionary theory predicts that the virulence and replication rates of these lineages should be lower than those of the lineages in mid-Central and East Africa. Seropositivity should also increase more gradually with age, more like the age structure of HIV-2 infections than pandemic HIV-1 infections (2, 3).

Paul W. Ewald
*Department of Biology,
Amherst College,
Amherst, MA 01002-2237*

REFERENCES

1. P. W. Ewald, *Hum. Nat.* 2, 1 (1991).
2. P. J. Kanki, R. G. Marlink, T. Siby, M. Essex, S. M'Boup, in *Advances in Applied Biotechnology Series*, vol. 7, *Gene Regulation and AIDS: Transcriptional Activation, Retroviruses, and Pathogenesis*, T. S. Papas, Ed. (Gulf, Houston, TX, 1990), pp. 255–272.
3. I. Romieu, R. Marlink, P. Kanki, S. M'Boup, M. Essex, *J. Acquired Immune Defic. Syndrome* 3, 220 (1990).
4. A. Del Mistro *et al.*, *ibid.* 5, 19 (1991).

Bevalac Funding

More is at stake in the imminent closure of Berkeley's Bevalac than David P. Hamilton suggests in his article "NASA researchers

protest DOE turnoff" (News & Comment, 5 June, p. 1388). For more than 15 years, heavy ion medical studies at the Bevalac have been unmatched throughout the world in studying the impact of using heavy charged particles to treat human cancers. No other facility currently in use can provide the wide range of beams in sufficient intensities for these studies, and the National Cancer Institute (NCI) has therefore provided support for clinical trials to determine the role of these beams in cancer therapy. The human trials, by their nature, are long-term studies and are not yet finished. The program recently underwent a competitive peer review as an NCI Program Project and obtained an excellent priority score that assures funding for the next 5 years, assuming continued beam time.

If the accelerator were shut down now, the phase II and III trials using high-energy helium and neon ions on various promising tumor sites would have to be halted before adequate statistics have been accumulated, and the United States would lose its leadership in the field of heavy ion studies. More important, patients for whom this therapy is the best hope of cure would be denied access to the beams. The next available facility is now being constructed in Japan and will not provide beams before 1995.

In addition, a newly designed (NCI-funded) beam scanning system for conformal therapy has just been built and tested, and the group is ready to begin the first human cancer patient treatments in the world with this new, precise therapy system. It will be possible with the new scanner to initiate true three-dimensional treatment planning and optimization with the use of these well-localizable charged particle beams. Also in jeopardy is a successful radiosurgical program funded by the Department of Energy for treating large arteriovenous malformations in the brain.

In short, the biomedical community as well as the NASA life science and instrumentation communities stand to lose substantially if the accelerator is closed prematurely. We urge that the decision for closure be reevaluated and that a unique and vital resource for U.S. and interested researchers abroad not be lost.

Joseph R. Castro

Department of Radiation Oncology,
University of California,
San Francisco, CA 94143, and
Lawrence Berkeley Laboratory,
Berkeley, CA 94720

Stanley B. Curtis

William T. Chu

Life Sciences Division,
Lawrence Berkeley Laboratory,
Berkeley, CA 94720

Missing Carbon Dioxide

Richard A. Kerr, in his Research News article "Fugitive carbon dioxide: It's not hiding in the ocean" (3 Apr., p. 35), wonders where about 3 to 4 billion tons of carbon emitted into the atmosphere each year are hiding, as they are unaccounted for in the current carbon budget. An article in the same issue by P. D. Quay *et al.* (p. 74) estimates the oceans are taking up about 2 billion tons of this, although Pieter Tans, as reported by Kerr, argues that it is closer to 1 billion tons a year. Research presented at a recent workshop cosponsored by the Environmental Protection Agency, Department of Energy, Edison Electric Institute, and the Umweltbundesamt (1, 2) concluded that a vast number of natural and managed ecosystems are sequestering excess carbon from fossil fuel and other sources in large enough amounts to readily account for the so-called "missing carbon."

What has stymied past attempts to balance the carbon budget is the unrealistic assumption that exchanges of CO₂ between terrestrial systems and the atmosphere are in steady state. That is, unless land is being converted from its natural state to a lower biomass (or carbon) system, the remainder of the landscape has a net CO₂ exchange of zero. This further implies that the whole natural landscape has not been disturbed by humans, which is, of course, unrealistic. The workshop participants concluded (2) that "New research, particularly from tropical forests, demonstrates that exchanges of CO₂ between the biota and the atmosphere have not been in steady state for several centuries." As these systems recover from past disturbances they can sequester substantial amounts of excess carbon from the atmosphere. Preliminary estimates of actual and potential carbon sinks in terrestrial and coastal systems were 3.7 to 9.1 billion tons of carbon a year, without consideration of CO₂ enrichment. More than 50% of these sinks are actual (in growth of temperate forests, coastal systems, soils, and some parts of the tropics); the rest are potential sinks that require new management regimes. Carbon accumulation in temperate forests has been confirmed by P. E. Kauppi *et al.* (Articles, 3 Apr., p. 70) as forests in Europe recover and accrete biomass from past disturbance.

Much of the research related to the tropics suggests that they are significant sources of atmospheric carbon. These estimates originate from models that are concerned mostly with the about 1 to 1.5% of forests that are being cleared every year; they assume that most of the other 98.5 to 99% or so of forest lands are in carbon steady state. The net flux could be consid-

erably smaller or even act as a sink (3). We know that large areas of these forests are undergoing gradual reductions in carbon stocks (4), but other areas are recovering—even the so-called mature forests—and are increasing their carbon stocks (3). This carbon can accumulate for several hundred years in large trees, soil organic matter, and woody debris.

Concerning the ability of ecosystems to continue to store carbon, participants of the workshop found that "Rising CO₂ accelerates net primary productivity and probably increases carbon storage in terrestrial ecosystems" (2). There is still plenty of "room" for expanding carbon storages (3), but the landscape has to be managed with strategies that enhance net carbon storage and are compatible with the conservation of biodiversity, sustainable land use, energy conservation, and economic development.

Sandra Brown

Department of Forestry,
University of Illinois,
Urbana, IL 61801

Ariel E. Lugo

Institute of Tropical Forestry,
Forest Service,

U.S. Department of Agriculture,
Southern Forest Experimental Station,
Río Piedras, Puerto Rico 00928

Joe Wisniewski

Wisniewski & Associates, Inc.,
6862 McLean Province Circle,
Falls Church, VA 22043

REFERENCES

1. J. Wisniewski and A. E. Lugo, Eds., *Water Air Soil Pollut.* 64 (nos. 1 and 2) (1992).
2. ———, *ibid.*, p. 455.
3. A. E. Lugo and S. Brown, *Forest Ecol. Manage.*, in press.
4. S. Brown, A. J. R. Gillespie, A. E. Lugo, *Can. J. Forest. Res.* 21, 111 (1991); S. Brown and A. E. Lugo, *Interciencia* 17, 8 (1992).

Malignant Hyperthermia

As David H. MacLennan and Michael S. Phillips correctly state in their article "Malignant hyperthermia" (8 May, p. 789), DNA-based testing may permit the preoperative identification of many individuals susceptible to malignant hyperthermia (MH). Contrary to a statement in Daniel E. Koshland, Jr.'s 8 May editorial (p. 717), however, knowledge of the molecular biology of MH has not yet reduced MH morbidity.

Current morbidity and mortality rates are unknown, but the dramatic decreases during the past decade are attributable to the efforts of (i) pharmacologists, medicinal chemists, and anesthesiologists who developed and tested dantrolene (1), the clinical