

Can afford.

Making Manual Plasmid Mini-Preps a Thing of the Past.

- **High Purity**
Sufficient for automated
sequencing and manual
transformations

Direct Pathways to Immunological Research

Taconic Transgenic Models

Transgenic technology offers gene-specific pathways to researchers for *in vivo* immunological research

pfp Knockout Mice
(Perforin Deficient)
severe depletion of
NK cell function

B2m Knockout Mice
(MHC Class I Deficient)
deficient in CD4⁺ T cells

AI Knockout Mice
(MHC Class II Deficient)
severe depletion of
CD4⁺ T cells

Rag-2 Knockout Mice
(Mature T and B
Cell Deficient)
unable to initiate
V(D)J rearrangement

TO ORDER

or for technical information:
e-mail: TaconicSer@aol.com
Tel: (518) 537-6208
Fax: (518) 537-7287

Taconic

Transgenic Models and Services Division

273 Haver Avenue
Germantown, NY 12526

fluence of carnivores in the site is given by five fossil remains of felids and mustelids (four individuals were adults). However, the low incidence of tooth marks (4% of the bone remains) especially in the joint attachments is not consistent with a pattern of carnivore priority access to the carcasses (2). On the contrary, the fossil assemblage of TD6 shows frequent dismembering and defleshing cutting marks on herbivore bone remains (7%). Further, most of the bone fragments show fractures produced when bones were still fresh, some of them with hammerstone impact notches likely produced during marrow extraction.

The bone assemblage from TD6 is consistent with butchering processes, and the presence of human remains at this site indicates conspecific consumption of human tissues.

Yolanda Fernández-Jalvo
J. Carlos Díez

José M. Bermúdez de Castro
Departamento Paleobiología,
Museo Nacional de Ciencias
Naturales (CSIC),
José Gutiérrez Abascal, 2,
28006 Madrid, Spain

Eudald Carbonell
Departamento Geografía e Historia,
Universitat Rovira i Virgili,
Plaza Imperial Tàrraco, 1,
43005 Tarragona, Spain

Juan Luis Arsuaga
Departamento Paleontología,
Universidad Complutense,
Ciudad Universitaria s/n,
28040 Madrid, Spain

References

1. P. Villa *et al.*, *Gallia Pr. Historie* **29**, 143 (1986a).
2. R. J. Blumenschine, *J. Archaeol. Sci.* **15**, 483 (1988); P. R. K. Richardson, *Palaeont. Afr.* **23**, 109 (1980).
3. J. Wymman, *Seventh Annual Report of the Trustees of the Peabody Museum of American Archaeology and Ethnology* **7** (1874); A. N. Somers, *Pop. Sci. Month.* **42**, 203 (1892); P. Villa *et al.*, *Science* **233**, 431 (1986); T. D. White, *Prehistoric Cannibalism at Mancos 5MTUMR-2346* (Princeton Univ. Press, Princeton, NJ, 1992), p. 462.
4. We are grateful to the SEM unit and the Restoration Laboratory of the Museo Nacional de Ciencias Naturales. Funds are provided by the Junta de Castilla y León and the Consejo Superior de Investigaciones Científicas.

Extracellular Protein Kinases

While viewing a poster on the involvement of ectoprotein kinase in the maintenance of hippocampal long-term potentiation during the recent Annual Meeting of the Society for Neuroscience (1), many of our colleagues pointed out that they have been misled by a statement in the Research News

article "A new face for the glutamate receptor" by Marcia Barinaga (13 Jan. 1995, p. 177). On page 178, this article states that there is evidence suggesting that "the big loop [in the glutamate receptor] is phosphorylated, and as phosphorylating enzymes act only within the cell, that would mean it must be inside the cell." The reader is left with the impression that the statement *phosphorylating enzymes act only within the cell* is an accepted scientific fact, but it contradicts over 50 scientific papers published on the activity of extracellular protein kinases and extracellular phosphoprotein phosphatases in various cell types. Most of these papers have appeared in prestigious journals. Several research groups have provided evidence that ectoprotein kinases operate on the cell surface, and a review article updated through 1990 is available (2).

In particular, with regard to receptor function, readers might be interested to learn that ectoprotein kinases are active on the surface of cells that store the co-substrate of protein kinase, adenosine triphosphate, within secretory vesicles and release it by exocytosis upon the stimulation of receptors or ion channels, such as platelets and neurons (3). Furthermore, the recent identification and characterization of an ectoprotein kinase with catalytic properties of atypical protein kinase C on the surface of brain neurons (1, 4) can shed new light on the controversy regarding the phosphorylation of extracellular sites in the glutamate receptors, and the potential involvement of such extracellular phosphorylation in long-term potentiation, a process implicated in the formation of memory traces in the brain.

Finally, clinical investigators also could benefit from learning about this research area, as exemplified by recent reports implicating ectoprotein kinase activity in the action of Alzheimer's amyloid peptides (4) and in malignant transformation (5).

Yigal H. Ehrlich

CSI/IBR, Center for Developmental Neuroscience,
College of Staten Island (CSI),
City University of New York,
and New York State Institute for
Basic Research in Developmental
Disabilities (IBR),
2800 Victory Boulevard,
Staten Island, NY 10314, USA

References

1. Y. H. Ehrlich *et al.*, *Neuroscience Abstr.* **21**, 598 (1995).
2. Y. H. Ehrlich, M. V. Hogan, Z. Pawlowska, U. Nalk, E. Kornecki, *Ann. N.Y. Acad. Sci.* **603**, 401 (1990).
3. Y. H. Ehrlich, T. B. Davis, E. Bock, E. Kornecki, R. H. Lenox, *Nature* **320**, 67 (1986); U. P. Nalk, E. Kornecki, Y. H. Ehrlich, *Biochim. Biophys. Acta* **1092**, 256 (1991).
4. M. V. Hogan, Z. Pawlowska, H.-A. Yang, E. Kornecki, Y. H. Ehrlich, *J. Neurochem.* **65**, 2022 (1995).
5. I. Friedberg, I. Belzer, O. Oged-Plesz, D. Kubler, *J. Biol. Chem.* **270**, 20560 (1995).

Downsizing at DOE Facilities

I would like to clarify a statement attributed to me in the News & Comment article "Deep cuts put heat on fusion, labs," by Andrew Lawler (3 Nov., p. 728). In a description of the impact of federal budget cuts, the National Institute for Petroleum and Energy Research (NIPER) in Bartlesville, Oklahoma, and two fossil fuel technology centers in West Virginia and Pennsylvania were listed as "likely targets for privatization."

The U.S. Department of Energy (DOE) is considering privatizing NIPER but we are not considering privatizing the Pittsburgh Energy Technology Center, or the Morgantown, West Virginia, Energy Technology Center. What we are doing is consolidating some management functions as part of DOE's restructuring and downsizing. As a result, the two facilities in West Virginia and Pennsylvania will operate under a single management structure. This consolidation will result in a reduction of some 90 positions, in total, at the two sites.

Dan W. Reicher

Acting Assistant Secretary for Policy,
U.S. Department of Energy,
Washington, DC 20585, USA


Comet Hunting: Discipline and Serendipity

The News article "Is Hale-Bopp the next great comet?" by Richard Kerr (29 Sep., p. 1818) describes the excitement felt by the astronomical community in anticipation of possibly the brightest comet of this century passing closest to the Earth. However, the article incorrectly identifies Alan Hale as an "amateur" astronomer. Hale received a Ph.D. in astronomy from New Mexico State University in 1993. Although Hale's primary research is not comets, he enjoys "comet-hunting" as a hobby. It was during one such session using his personal telescope in his backyard last July that he stumbled upon what may be the great comet of the 20th century. Serendipity continues to play an important role in astronomical discovery.

Jack O. Burns

Department of Astronomy,
New Mexico State University,
Las Cruces, NM 88003, USA
E-mail: jburns@nmsu.edu

Phosphocellulose Paper In a 96-Well Plate



Increase Throughput For Kinase Assays

Signal transduction assays are an important tool in drug discovery. The Millipore MultiScreen Assay System uniquely offers a 96-well filter plate containing phosphocellulose paper and protocols to perform peptide substrate kinase assays. No more cutting and transferring the paper! Every step, from the phosphorylation reaction to direct scintillation counting, can be carried out in one plate.

The MultiScreen Assay System can also be used with other filter formats, including glass fiber or low protein binding Durapore® (PVDF) membrane – both ideally suited for TCA precipitated substrates.

Call or fax for more information. U.S. and Canada, call Technical Services: 1-800-MILLIPORE (645-5476); in Japan, call: (03) 3474-9111; in Europe, fax: +33.88.38.91.95.

MILLIPORE

MILLIPORE LAB CATALOG ON INTERNET:
ACCESS URL MENU AND TYPE:
<http://www.millipore.com>