items stored in -70° C freezers, but the cycling did not maintain desired temperatures in -20° C freezers or in refrigerators. Also, for a time the building was inaccessible or could not be entered except by maintenance personnel, and so stored items could not be moved (and there was essentially nowhere in town to move them to, in any case). The result was that five floors of research laboratories lost numerous heat-labile enzymes and other reagents because they did not survive the temperature fluctuations.

After the flooding, four companies offered significant levels of aid to researchers in Grand Forks. First, Strategene rapidly contacted departments in the School of Medicine and offered to resupply, gratis, all enzymes and reagents that were lost by our laboratories as a result of the flood, regardless of their original commercial source. This saved a huge amount of time and funding for several laboratories, including my own and three others in my department. In addition, Stratagene made available to us additional materials and equipment at a large discount. Boehringer Mannheim Biochemicals resupplied several laboratories with reagents, also at a huge discount, and Glaxo Wellcome sent many items of equipment to either replace or enhance specific research capabilities damaged during the flooding. Finally, Beckman Instruments donated to the Department of Microbiology and to the Department of Biochemistry several microcentrifuges, clinical centrifuges, pH meters, and spectrophotometers to equip the undergraduate and medical teaching laboratories that had been destroyed.

The actions of these companies significantly reduced our research and teaching downtime during a period when all of us were occupied with conflicting demands in reconstructing our community and our personal, professional, and teaching lives. Each company approached us unasked, and deserves our public thanks for their generosity.

Kevin D. Young Thomas Hill Ann Flower Francis Sailer Department of Microbiology and Immunology, University of North Dakota School of Medicine, Grand Forks, ND, USA

Biodiversity and Ecosystem Properties

Because the goal of ecology is to understand the causes of patterns observed in the natural world, studies that document novel patterns in natural ecosystems are of central importance. The report "The influence of island area on ecosystem properties" by David A. Wardle et al. (29 Aug., p. 1296) is such a study. Some interpretations of data made by Wardle et al. illustrate the difficulty of ascribing causation on the basis of observation. In particular, their assertions "that ecosystem process rates were lowest on those islands with the greatest diversity" and that "this finding is in direct contrast to other studies which have shown elevated process rates in more diverse communities" are difficult to justify given their analyses and the data in the other studies (1-4).

In the archipelago studied by Wardle *et al.*, the variables of island size, fire frequency, successional state, and species composition were strongly interdependent. Compared with larger islands, smaller islands were found to have lower fire frequencies, higher plant diversity, and species compositions biased toward later successional species. Because of these interdependencies, correlations between any one of these variables and any of the ecosystem response variables may lead one to spurious conclusions. Multiple regressions, or other multivariate techniques, would be needed to determine if the data

"These columns add just the right spin to my lab work: they get it done so that I can have fun," says Marc Awobuluyi, a molecular neurobiologist living and working in Boston, Massachusetts, USA.

What do I need to get Hi purity of a biomolecule with a single Trap step

(The answer is in the question, Marc) LETTERS

support the hypothesis that a given ecosystem response variable significantly depends on diversity, or composition, or fire frequency. Once there has been statistical control for correlations among these variables. Even when done carefully, such analyses of patterns in natural systems are still open to alternative interpretations.

Such difficulties have caused ecology to become an increasingly experimental science. Direct experimental tests of the effects of plant diversity on ecosystem properties (1-4) may give more valid results than would correlations that are uncorrected for collinear variables. The most significant advances come when results of experiments, observational studies, and theory (5) are congruent. Wardle et al. do not mention some aspects of these diversity experiments. All have found that plant species diversity or plant functional diversity, or both, have significant effects on ecosystem variables, but some of these rates were highest (1-3) and some were lowest (3, 4) in habitats with high diversity, and some seemed to vary idiosyncratically with diversity (1). In particular, there were lower soil concentrations of nitrate and ammonium in habitats with higher diversity (3, 4), which, contrary to statements by Wardle et al., is consistent with the correlation they report.

We commend Wardle et al. on identifying a natural system that may provide significant insights into the forces controlling the functioning of ecosystems, but suggest that more complex analyses and experimental manipulations are needed to obtain the deeper insights that this system may offer.

> David Tilman Shahid Naeem Johannes Knops Peter Reich Evan Siemann Department of Ecology, Evolution and Behavior, University of Minnesota, St. Paul, MN 55108-6097, USA E-mail: tilman@lter.umn.edu David Wedin Department of Botany, University of Toronto, Toronto, Ontario, M5S 3B2, Canada Mark Ritchie Department of Fisheries and Wildlife, Utah State University, Logan, Utah 84322-5210, USA John Lawton Centre for Population Biology, Imperial College at Silwood Park, Ascot, Berks, SL5 7PY, United Kingdom

References

- 1. S. Naeem, L. J. Thompson, S. P. Lawler, J. H. Lawton, R. M. Woodfin, Nature 368, 734 (1994). 2. S. Naeem, K. Håkenson, J. H. Lawton, M. J. Craw-
- ley, L. J. Thompson, Oikos 76, 259 (1996). 3. D. Tilman, D. Wedin, J. Knops, Nature 379, 718
- (1996); D. Tilman, J. Knops, D. Wedin, P. Reich, M. Ritchie, E. Siemann, Science 277, 1300 (1997). 4. D. U. Hooper and P. M. Vitousek, Science 277, 1302
- (1997). 5. D. Tilman, C. L. Lehman, K. T. Thomson, Proc. Natl.
- Acad. Sci. U.S.A. 24 1857 (1997).

Response: The island system that we considered included islands with high plant diversity and those with low diversity. If plant diversity was an important causative agent for maintaining ecosystem properties in nature, then we would expect measurements of these properties to be greatest on islands with the greatest diversity. This was not the case, whether or not we analyzed the data by multivariate and multiple regression approaches correcting for the effects of island area and species composition. Although the patterns that we observed cannot be explained in terms of diversity effects, they can in terms of plant compositional effects and in particular the ecophysiological traits of the dominant plant species present. Our results are consistent with studies that found (i) little evidence in nature for consistent positive relationships between spe-



Which biomolecule have you been working with lately? A nucleic acid? A peptide? A protein? Whichever biomolecule you work with, you'll need to capture and purify it to move ahead in your work. Small, pre-packed chromatography cartridges are the easiest one-step solution for quickly moving all the way from scouting to large scale purification. These products are called HiTrap".

HiTrap: best for easy one-step purification

HiTrap is simply the most versatile range of easy-to-use pre-packed cartridges ever available for one-step purification. Individual HiTrap products provide just what you need to master a wide variety of applications—including the isolation and purification of: His-tagged fusion proteins and peptides, nucleotiderequiring enzymes, monoclonal antibodies, polyclonal antibodies, DNA-binding proteins, and more.

You can choose the method of purification that suits you best as any individual HiTrap can be used with a syringe, a pump or a system. As to their performance, HiTrap recombinant protein A (pictured; top left) has a 30-40% higher dynamic binding capacity when compared to similar products.

Find out more about the HiTrap range. Give us a call: 1 (800) 526-3593 in the USA; +81 3492 6949 in Japan; +46 18 16 50 11 in Europe and the rest of the world.

Or visit us on the Internet: http://www.biotech.pharmacia.se.



- Exclusively for biomedical research at universities and nonprofit scientific institutions.
- The laboratory has studied normal and abnormal development and provided tissue for 35 years.
- Most tissues are available for study.
- Tissues can be supplied from most gestational stages and from normal or abnormal specimens.
- Immediate processing includes rapid fixation, LN₂, balanced salt or medium as requested.
- Tissues shipped nationwide by overnight air are suitable for molecular biology, enzymology, receptor study, electron microscopy, etc.

For further information:

Phone 800-583-0671 FAX 800-583-0668

Please include an abstract of proposed research, preferably on PHS 398. Circle No. 23 on Readers' Service Card

cies diversity and ecosystem properties (1), (ii) idiosyncratic relationships between plant diversity and the functioning of decomposition-related processes (2), and (iii) plant physiological attributes to be important in determining ecosystem-level properties (3).

Tilman et al. state that we "do not mention some aspects" of previous diversity experiments and cite several references (4-7), but two of those (4) were not published until after our report was accepted. The other studies (5-7) found greater rates of processes in the highest diversity treatments for the majority of ecosystem properties that showed significant treatment responses. It is presumably those findings that induced Tilman et al. (5, p. 720) to conclude that reduced biodiversity "threaten ecosystem productivity and the sustainability of nutrient cycling" and Naeem et al. (6, p. 735) to state that loss of biodiversity will "alter or impair the services that ecosystems provide" [italics ours]. Our data are not in agreement with those findings, which are based on experiments in which the most diverse treatments may have a greater probability of being dominated by the most productive species in the entire species pool (8). Our results are in much closer agreement with recent studies (4) which are designed more robustly and are less likely to be influenced by this problem; these studies conclude that species composition, rather than species diversity, is the main determinant of ecosystem properties.

The data about mineral N presented by Tilman et al. (5) are not comparable with our data. In their study, the reduction of soil mineral N in the most species-diverse plots was presumably a result of the enhanced rate of a key process (plant productivity) that led to more complete N uptake. In our study, accumulation of organic N in the soil of the most species-diverse islands was induced by retardation of key soil processes and resulted in long-term lock-up of N in biologically inaccessible pools.

We agree with Tilman et al. about the utility of experimental and theoretical approaches and in this light have recently established long-term manipulative experiments on our island system. One experiment, initiated in 1996, aims to determine the ecosystem-level significance of both plant species and functional group diversity across 30 of our islands; it includes the use of 420 experimental plots. We believe, however, that experimental and theoretical approaches are only a means to an end (the end being to better understand what happens in real ecosystems) rather than an end

Redefining One-Step Bibliographies

NEW—Reference Manager. V8 for Windows 95/NT



"I have been using Reference Manager for 10 years and now can't imagine writing a manuscript or grant without it. The new Cite While You Write feature is a great addition to the program and makes the process of preparing manuscripts even easier than before.'

> Grav F. Crouse Department of Biology Emory University

Easy One-Step Bibliographies

- · Generate your manuscript and bibliography as a single document - in one step
- Never leave your word processor to identify citations
- Create in-text citations as author/date or numbered
- · Format bibliographies for hundreds of journal styles, or create your own
- Revise your manuscript as many times as needed even make complete style changes

Exclusive features

- Perform key operations across multiple databases
- Link directly to the Web
- Store electronic citations & Internet communications in ready to use templates
- · Create synonyms for periodicals, keywords and authors
- Unlimited database capacity & number of databases

NEW — true multi-user network solutions!

AND MUCH MORE SPECIAL PRICE FOR SCIENCE SUBSCRIBERS

Reference Manager and Capture \$199.95 through December 31, 1997 (single-user version 30-Day Money-Back Guarantee)

RIS

Research Information Systems, 2355 Camino Vida Roble, Carlsbad, CA 92009 USA 800.722.1227 • 760.438.5526 • Fax: 760.438.5266 • E-mail: sales@risinc.com • WWW: http://www.risinc.com

Circle No. 33 on Readers' Service Card

Division of Institute for Scientific Information A Thomson Company Letners

in themselves. We are not persuaded that experimental tests should have precedence, and believe that if outcomes of short-term experimental (5, 6) and theoretical studies do not concur with patterns and processes observed in nature (1 and our report), then it is the experiments and theory that should be queried.

David A. Wardle Landcare Research, Post Office Box 69, Lincoln 8152, New Zealand Olle Zackrisson Greger Hörnberg Swedish University of Agricultural Sciences, S901-83 Umeå, Sweden Christiane Gallet Université de Savoie, F-73376 Le Bourget-du-Lac Cedex, France

References

- M. M. Al Mufti, C. L. Sydes, S. B. Furness, J. P. Grime, S. R. Band, *J. Ecol.* 65, 759 (1977); J. P. Grime, *Plant Strategies and Vegetation Processes* (Wiley, Chichester, UK, 1979); G. M. Rusch and M. Oesterheld, *Oikos* 78, 519 (1997).
- K. Chapman, J. B. Whittaker, O. W. Heal, Agric. Ecosyst. Environ. 24, 33 (1988); D. A. Wardle, K. I. Bonner, K. S. Nicholson, Oikos 79, 247 (1997).
- J. H. C. Cornelissen and K. Thompson, *New Phytol.* 135, 109 (1997); D. A. Wardle, G. M. Barker, K. I. Bonner, K. S. Nicholson, *J. Ecol.*, in press.
- D. U. Hooper and P. M. Vitousek, *Science* 277, 1302 (1997); D. Tilman, J. Knops, D. Wedin, P. Reich, M. Ritchie, E. Siemann, *ibid.*, p. 1300.
 D. Tilman, D. Wedin, J. Knops, *Nature* 379, 718
- 5. D. Tilman, D. Wedin, J. Knops, *Nature* **379**, 718 (1996).
- S. Naeem, L. J. Thompson, S. P. Lawler, J. H. Lawton, R. M. Woodfin, *ibid.* **368**, 734 (1994).
- S. Naeem, K. Håkenson, J. H. Lawton, M. J. Crawley, L. J. Thompson, *Oikos* **76**, 259 (1996).
- 8. M. A. Huston, Oecologia 110, 449 (1997).

Corrections and Clarifications

In the Perspective "Do continents part passively, or do they need a shove?" by R. W. Carlson (10 Oct., p. 240), figure 1, with a caption reading, "Flood basalt. A 2-km-thick section of the Columbia River basalts in the Grande Ronde River valley of eastern Washington," is in fact a view of the Kiger River Gorge in the northern part of Steens Mountain in southeastern Oregon.

Letters to the Editor

Letters may be submitted by e-mail (at science_letters@aaas.org), fax (202-789-4669), or regular mail (*Science*, 1200 New York Avenue, NW, Washington, DC 20005, USA). Letters are not routinely acknowledged. Full addresses, signatures, and daytime phone numbers should be included. Letters should be brief (300 words or less) and may be edited for reasons of clarity or space. They may appear in print and/or on the World Wide Web. Letter writers are not consulted before publication.



Note to Readers about Technical Comments

Beginning in 1998, summaries of technical comments will appear in the printed journal; full text and figures will appear only online.

This change will allow:

■ **Free access** to the full text of technical comments and responses for all *Science* Online readers.

■ Quicker publication of reviewed comments, criticisms, and responses.

■ Linkage between reports and comments, which would notify readers that subsequent comments were published about a report (or article) in *Science*.

■ More comprehensive figures and tables in the comments and more detailed discussion of methods and instrumentation.

■ **Up-front coverage** in summaries of the comments and responses, which will appear in "This Week in Science" in the printed journal, with Web addresses for the full texts.

■ Continued full citation, with technical comment titles and authors continuing to be listed in the printed Table of Contents, and authors continuing to be cited in the quarterly online index.

TIME MACHINE



- **Fast**—up to 24 preps per hr, saving you valuable time.
- High Purity—sufficient for automated fluorescent and manual sequencing.
- Easy Operation—begin prep with direct loading of bacterial culture. No centrifugation step saves you time.
- **Consistent Results**—up to 6 µg of plasmid per ml. Quality DNA … time and time again.

Call now to learn how the New Improved Mini-Prep 24 can give you quality DNA and save time by automating your Plasmid DNA preps!



¹¹³³⁹ Sorrento Valley Road San Diego, CA 92121 (619) 452–2603 • Fax (619) 452–6753 www.macconnell.com Circle No. 29 on Readers' Service Card