

Modulation of Respiratory Frequency by Peptidergic Input to Rhythmogenic Neurons in the PreBötzinger Complex

Paul A. Gray,^{1,3*} Jens C. Rekling,^{1*} Christopher M. Bocchiaro,²
Jack L. Feldman^{1,2†}

Neurokinin-1 receptor (NK1R) and μ -opioid receptor (μ OR) agonists affected respiratory rhythm when injected directly into the preBötzinger Complex (preBötC), the hypothesized site for respiratory rhythmogenesis in mammals. These effects were mediated by actions on preBötC rhythmogenic neurons. The distribution of NK1R⁺ neurons anatomically defined the preBötC. Type 1 neurons in the preBötC, which have rhythmogenic properties, expressed both NK1Rs and μ ORs, whereas type 2 neurons expressed only NK1Rs. These findings suggest that the preBötC is a definable anatomic structure with unique physiological function and that a subpopulation of neurons expressing both NK1Rs and μ ORs generate respiratory rhythm and modulate respiratory frequency.

Respiratory rhythm is generated by neurons in the brainstem. Surprisingly, given the vital role and the apparent simplicity of breathing movements, the identity of these neurons and the mechanisms of respiratory rhythmogenesis and frequency modulation are unknown. Respiratory neurons are found in a narrow column in the ventrolateral medulla extending from the facial nucleus to the spinal cord (1, 2). The preBötC is the limited portion of this column necessary and sufficient for in vitro neonatal rodent medullary slice preparations to generate respiratory-related motor nerve output (2, 3). Perturbations of neuronal function within the preBötC in adult rats and cats severely disrupt breathing (4), consistent with the hypothesis that this region contains the neurons responsible for generating respiratory rhythm. Unfortunately, these and related experiments only circumscribe the approximate boundaries of the preBötC. In the absence of a precise anatomic delineation of the preBötC, defining its functional role or the roles of individual neurons it contains is problematic.

Local application of the NK1R agonist, substance P (SP), into the preBötC in vitro increases the frequency of endogenous respiratory-related output; similar application of the μ OR agonist [D-Ala², N-Me-Phe⁴, Gly⁵-ol]enkephalin acetate (DAMGO) or the γ -aminobutyric acid type B receptor (GABA_BR) agonist baclofen

decreases respiratory frequency (5–7). Similar responses are seen in vivo (8). Because these drugs affect frequency, they must act either directly on preBötC rhythmogenic neurons or indirectly on other preBötC neurons that provide modulatory signals to the rhythm generator. In vitro respiratory-related output does not require chlorine-mediated postsynaptic inhibition (2, 3, 5, 9). If peptide neuromodulators affect respiratory frequency by direct actions on only inhibitory neurons, as has been suggested for opioid actions in the midbrain (10), then peptide receptor-expressing neurons could not be responsible for rhythm generation. Thus, we examined whether the effects of SP and DAMGO on respiratory rhythm in vitro were affected by blocking GABA type A receptor (GABA_AR) and glycine receptor activation (11). Pressure injection of SP (2 pmol) into the preBötC of rhythmically active brainstem slices from rats increased respiratory frequency to $192\% \pm 5\%$ of control, whereas injection of DAMGO (1.5 pmol) decreased frequency to $64\% \pm 1\%$ of control (Fig. 1). GABA_BR activation has similar effects (5, 7). Bath application of the GABA_AR antagonist bicuculline and the glycine receptor antagonist strychnine to these slices did not change the baseline respiratory frequency or the effects on respiratory frequency of SP or DAMGO (Fig. 1B). Thus, the preBötC NK1R-, GABA_BR-, and μ OR-expressing neurons that mediate agonist frequency-modulating effects are likely excitatory and are either directly involved in respiratory rhythm generation or are nonrespiratory neurons that modulate rhythm-generating circuits.

We examined the immunohistochemical expression pattern of NK1R, μ OR, and GABA_BR in the ventrolateral medulla of

adult rats (12). Within the ventrolateral respiratory column, GABA_BR expression was widespread. However, both NK1R and μ OR expression were concentrated within the preBötC and absent from the adjacent Bötzinger complex (BötC) (1) and rostral ventral respiratory group (1) (Fig. 2); NK1R expression in adult mouse brainstem was similar (13). The preBötC NK1R⁺ neurons were segregated from both motoneurons and catecholaminergic sympathetic neurons as revealed by double immunohistochemistry for choline acetyltransferase (ChAT) or tyrosine hydroxylase (TH) (13, 14).

To determine whether GABAergic and peptidergic effects on breathing are due to convergent inputs onto individual neurons, we looked for μ OR or GABA_BR colocalization in NK1R⁺ neurons. We found NK1R⁺/ μ OR⁺ neurons and processes (Fig. 3A) and NK1R⁺/ μ OR⁺ neurons and processes (of 88 preBötC NK1R⁺ neurons examined in detail,




Fig. 1. Effects of peptides injected into preBötC are independent of GABA_A and glycinergic transmission. (A) Microinjection into preBötC of DAMGO (2 pmol) slows and SP (1.5 pmol) speeds up endogenous respiratory-related rhythm. Traces are integrated XII motor nerve output; bars under each trace represent injection period with total amount indicated. (B) Frequency response to microinjection of DAMGO and SP into preBötC is not affected by blocking chlorine-mediated postsynaptic transmission with bath application of bicuculline (20 μ M) and strychnine (1 μ M) sufficient to block the effects of the combined microinjection of the GABA_A agonist, THIP (1 nmol), and glycine (1 nmol). Histograms of grouped data across multiple experiments are shown; number of epochs (20 cycles per epoch) are indicated within bars; error bars are SEM. Statistical comparisons are between individual pairs of epochs in the same experiment: *, significant difference ($P < 0.05$; Student's *t* test) between conditions in all pairs; ns, no significant difference between conditions in all pairs; §, significant difference ($P < 0.05$) relative to control in all pairs; ns, no significant difference relative to control in all pairs. Bic/Str, bath application of bicuculline and strychnine.

Departments of ¹Neurobiology and ²Physiological Science and ³Interdepartmental Ph.D. Program in Neuroscience, University of California Los Angeles, Box 951763, Los Angeles, CA 90095–1763, USA.

*These authors contributed equally to this work.
†To whom correspondence should be addressed. E-mail: feldman@ucla.edu

REPORTS

53 coexpressed μ OR) (Fig. 3B). We found $NK1R^+/GABA_B R^+$ (Fig. 3C) as well as $NK1R^+/GABA_B R^-$ neurons (of 160 preBötC $NK1R^+$ neurons examined in detail, 118 coexpressed $GABA_B R$). We also found $NK1R^-/GABA_B R^+$ neurons (Fig. 3C) and a few $\mu OR^+/NK1R^-$ processes (Fig. 3B) and somas (13). Thus, GABAergic and peptidergic synaptic inputs converge on subsets of preBötC neurons. Moreover, in the ventrolateral respiratory column, neurons that are targets of these convergent modulatory inputs that affect respiratory frequency are present only in the preBötC. We propose that the expression of $NK1R^+$ neuronal somata in the ventrolateral respiratory column defines the anatomic extent of the preBötC.




Fig. 2. $NK1R^-$ and μOR -expressing neurons are limited to and define the preBötC. (A) Cartoon indicating approximate location and boundaries of ventral respiratory column regions and adjacent ambigua motoneuron pools in sagittal section. (B) $NK1R$ immunohistochemical expression in a single sagittal section through the respiratory column corresponding to box in (A). Higher magnification images showing cell soma staining from preBötC (C) and relative lack of staining in BötC (D). Confocal, negative composite images of μOR immunohistochemical staining indicating different expression levels in the preBötC (E) (blue arrow) and BötC (F) (red arrow) in transverse section. Note high levels of expression of these peptide receptors in the cNA, a cranial motoneuron pool not related to respiration. Scale in micrometers. cNA, compact formation of the nucleus ambiguus; scNA, subcompact formation of NA; rVRG, rostral ventral respiratory group; VII, facial motor nucleus; D, dorsal; L, lateral.

These results do not differentiate whether these preBötC neurons generate or simply modulate respiratory rhythm. We tested the responsiveness of behaviorally identified respiratory neurons to neurokinin and opioid agonists (15). In the preBötC of rhythmically active slices from mice, inspiratory neurons fall into two categories (types 1 and 2) based on their electroresponsive properties and membrane potential trajectories (16, 17) (Fig. 4A). After blockade of action potential driven synaptic transmission by 2 μM tetrodotoxin (TTX), bath application of SP depolarized type 1 neurons 7.7 ± 1.5 mV ($n = 4$) (Fig. 4B), whereas type 2 neurons depolarized much less [2.7 ± 0.8 mV ($n = 4$) (Fig. 4B)]. Input resistance was not significantly changed by SP in type 1 or type 2 neurons (Wilcoxon signed rank test). Bath application of DAMGO (in TTX) hyperpolarized all type 1 neurons tested (7.9 ± 1.8 mV; $n = 5$) and reduced their input resistance to 67% of control value but had no effect on either the membrane potential or input resistance of type 2 neurons ($n = 6$) (Fig. 4C). The current-voltage ($I-V$) curves of type 1 neurons before and after DAMGO (under voltage clamp in the presence of TTX) intersected at -102 mV with normal $[K^+]_o$ concentration ($n = 5$), whereas when $[K^+]_o$ was increased by 10 mM, the intersection point shifted to -66 mV ($n = 3$; $P < 0.05$, Mann-Whitney rank sum test) (Fig. 4D). Taken together, these results suggest that the effect of DAMGO was mediated by activation of a K^+ conductance.




Fig. 3. G-protein-coupled receptors are coexpressed on individual preBötC neurons. Confocal projections of $NK1R^+$ somas (A to C) (left) and processes (B) (left), μOR^+ somas (A) (right) and processes (A and B) (right), and $GABA_B R^+$ somas (C) (right). Arrows indicate coexpression and arrowheads indicate absence of coexpression. Bars = 25 μm .

Most models for respiratory rhythm generation predict that depolarization of rhythm-generating neurons will speed up rhythm and that their hyperpolarization will slow it down (2, 3, 18). Type 1 neurons, which are rhythmogenic (2, 16, 17), are the only preBötC inspiratory neurons depolarized by both SP and thyrotropin-releasing hormone (16), peptides that increase respiratory frequency, and hyperpolarized by DAMGO, a peptide that decreases respiratory frequency; this is consistent with these models and the hypothesis (2) that type 1 neurons are responsible for both rhythmogenesis and frequency control.

The preBötC appears to have an important function in generating respiratory rhythm and in processing signals affecting respiratory frequency (2). We propose that the kernel for




Fig. 4. Postsynaptic membrane responses of inspiratory neurons in the preBötC to SP and DAMGO. (A) Characteristic membrane potential trajectory between inspiratory discharges of type 1 and type 2 inspiratory neurons. Note the long-lasting hyperpolarization characteristic of type 1 neurons. (B) Membrane responses of type 1 and type 2 neurons to bath-applied SP (1 μM) after blockade of action potential driven synaptic transmission by TTX (2 μM). Hyperpolarizing current pulses (1-s duration every 5 s) were injected to measure input resistance throughout the application and the membrane potential was brought back to control value by injecting bias current at the indicated time (bias I). (C) Membrane responses of type 1 and type 2 neurons to bath-applied 5 μM DAMGO (TTX). (D) (Top) $I-V$ curves (in the presence of TTX) before and after DAMGO application (5 μM) in normal $[K^+]_o$ (6.2 mM). Arrow indicates potential at which the two curves intersect. (Bottom) $I-V$ curves (in the presence of TTX) before and after DAMGO application (5 μM) in high $[K^+]_o$ (16.2 mM). Note intersection points of the curves have shifted in the depolarizing direction.

the basic respiratory rhythm is a small class of preBötC rhythmogenic neurons (type 1) onto which several modulatory systems converge. These neurons may overlap or even be identical to other preBötC neurons with intrinsic oscillatory bursting properties—for example, pace-maker neurons (3, 19) and preinspiratory neurons (20)—because 50% of type 1 neurons have bursting properties (16) and at least some project to the midline (21). However, whether endogenous burst activity is critical for generation of respiratory rhythm per se (3, 18–20) remains to be determined. Alterations in peptidergic transmission in the preBötC may play a role in respiratory disorders such as sleep apnea (22) and sudden infant death syndrome (23). The possibility that multiple peptide systems affecting respiration converge on a particular identifiable class of neurons represents an interesting locus for ventilatory control. The coexpression within the ventrolateral respiratory column of NK1R and μ OR in preBötC neurons provides an opportunity to exploit targeted molecular manipulation of breathing in animal models and a basis for analysis of human respiratory disorders.

References and Notes

- A. L. Bianchi, M. Denavit-Saubie, J. Champagnat, *Physiol. Rev.* **75**, 1 (1995); J. L. Feldman, in *Handbooks of Physiology: The Nervous System. Intrinsic Regulatory System in the Brain*, F. Bloom, Ed. (American Physiological Society, Bethesda, MD, 1986), pp. 463–524; E. G. Dobbins and J. L. Feldman, *J. Comp. Neurol.* **347**, 64 (1994).
- J. C. Reklung and J. L. Feldman, *Annu. Rev. Physiol.* **60**, 385 (1998).
- J. C. Smith, H. H. Ellenberger, K. Ballanyi, D. W. Richter, J. L. Feldman, *Science* **254**, 726 (1991).
- A. Monnier, F. Hayashi, D. R. McCrimmon, *Soc. Neurosci. Abstr.* **346**, 5 (1998); N. Koshiya and P. G. Guyenet, *J. Physiol. London* **491**, 859 (1996); J. M. Ramirez, S. W. Schwarzacher, O. Pierrefiche, B. M. Olivera, D. W. Richter, *J. Physiol. London* **507**, 895 (1998); J. H. Hsieh et al., *J. Auton. Nerv. Sys.* **73**, 7 (1998).
- J. L. Feldman and J. C. Smith, *Ann. NY Acad. Sci.* **563**, 114 (1989).
- S. M. Johnson, J. C. Smith, J. L. Feldman, *J. Appl. Physiol.* **80**, 2120 (1996).
- J. Brockhaus and K. Ballanyi, *Eur. J. Neurosci.* **10**, 3823 (1998).
- A. C. Bonham, *Resp. Physiol.* **101**, 219 (1995); D. R. McCrimmon and G. S. Mitchell, in *Regulation of Breathing*, J. A. Dempsey and A. I. Pack, Eds. (Dekker, New York, 1995), vol. 79, pp. 151–218.
- G. D. Funk, J. C. Smith, J. L. Feldman, *J. Neurophysiol.* **70**, 1497 (1993); X. M. Shao and J. L. Feldman, *J. Neurophysiol.* **77**, 1853 (1997).
- H. L. Fields and M. M. Heinricher, *Philos. Trans. R. Soc. London Ser. B* **308**, 361 (1985).
- Transverse 400- or 800- μ m-thick slices of the brainstem from BALB/c mice (P_0 – P_3) and Sprague-Dawley rats (P_0 – P_4) were cut at the level of the caudal and rostral preBötC, respectively, as reported (3) and perfused in artificial cerebrospinal fluid (aCSF) containing 128 mM NaCl, 9 mM KCl, 0.5 mM NaH_2PO_4 , 1.5 mM CaCl_2 , 1.0 mM MgSO_4 , 23.5 mM NaHCO_3 , 30 mM glucose (rat), or 130 mM NaCl, 5.4 mM KCl, 0.8 mM KH_2PO_4 , 26 mM NaHCO_3 , 30 mM glucose, 1 mM MgCl_2 , 0.8 mM CaCl_2 (mouse). Rhythmic respiratory-related motor output was recorded by a suction electrode placed on a XII nerve rootlet. Nerve activity was amplified, filtered at 3 Hz to 1 kHz, and rectified nerve activity was integrated. SP, DAMGO, 4,5,6,7-tetrahydroisoxazolo[5,4-c]pyridin-3-ol (THIP) hydrochloride, glycine, and TTX were obtained from RBI and dissolved in aCSF. Drugs were microinjected with glass micropipettes (tip diameter 6 to 12 μ m) lowered (100 to 300 μ m) into the preBötC by micromanipulator. Injections were made with a controlled pressure source for 5 s. Ejection volume was monitored with a calibrated eyepiece reticule.
- Female adult Sprague-Dawley rats ($n = 17$) and BALB/c mice ($n = 2$) were transcardially perfused with 4% paraformaldehyde in phosphate-buffered saline (PBS) with or without picric acid, cryoprotected in 25% sucrose PBS, embedded in OCT, sectioned at 25 to 40 μ m on a cryostat, and processed free floating. Sections were incubated in primary antibody diluted in serum at 4°C for 24 to 48 hours, placed in fluorescent (Jackson ImmunoResearch) or biotin (Vector Laboratories) conjugated species-specific secondary antibody overnight, and mounted on gelatin-subbed slides. In several rats, μ OR immunohistochemistry was amplified with tyramine signal amplification—direct green amplification (NEN LifeSciences) and bright-field NK1R staining was visualized with a Vectastain Elite ABC kit and diaminobenzidine. Antibodies: rabbit antibody to NK1R (anti-NK1R) (1:2500 fluorescent, 1:25,000 bright field; Chemicon), guinea pig anti-GABA_BR (1:3000; Chemicon), goat anti-CHAT (1:100; Chemicon), mouse anti-TH (1:1000; Boehringer Mannheim), rabbit anti- μ OR (1:75,000 TSA or 1:7000 fluorescent; Instar). Bright-field images were digitally acquired (Polaroid DMC) into Photoshop (Adobe Systems) and background subtracted to remove chip nonlinearities. Confocal images were acquired with a Zeiss 410 SCM and maximal intensity projections were generated in NIH Image. All images were filtered and adjusted for contrast and light levels in Photoshop for clarity. Cell counts for coexpression were obtained by visual inspection of confocal images ($n = 2$).
- P. A. Gray and J. L. Feldman, data not shown.
- J. L. Feldman and H. H. Ellenberger, *Annu. Rev. Physiol.* **50**, 593 (1988); H. H. Ellenberger, J. L. Feldman, W. Z. Zhan, *J. Comp. Neurol.* **294**, 212 (1990); H. H. Ellenberger and J. L. Feldman, *J. Comp. Neurol.* **294**, 202 (1990).
- Neurons located at the caudal or rostral part of the preBötC in mice were visualized with differential interference contrast and infrared video microscopy. Glass micropipettes (resistances typically were 2 to 3 MW) were filled with a solution containing 115 mM HMeSO_3 , 115 mM KOH, 5 mM NaCl, 1 mM MgCl_2 , 0.01 mM CaCl_2 , 0.1 mM BAPTA [1,2-bis(2-aminophenoxy)ethane- N,N,N',N' -tetraacetic acid, tetra- K^+ salt], 10 mM Hepes, 3 mM 2-ATP(Mg^{2+}) (pH 7.3). Current- or voltage-clamp recordings were made with an Axoclamp-2A (Axon Instruments) amplifier in bridge or single electrode voltage clamp mode. The I - V relationship was determined by injecting a slow (7-s cycle) ramp-shaped current and plotting the rising phase of the I - V . Signals were recorded on videocassette (pulse code modulation; Vetter Instruments, model 3000A), digitized at 1 to 20 kHz with a Digidata 1200 A/D board (Axon Instruments) and analyzed with Axoscope software (Axon Instruments). Statistical values are given as mean $\pm$ SEM.
- J. C. Reklung, J. Champagnat, M. Denavit-Saubie, *J. Neurophysiol.* **75**, 811 (1996).
- , *J. Neurophysiol.* **75**, 795 (1996).
- R. J. Butera Jr., J. Rinzel, J. C. Smith, *J. Neurophysiol.* **82**, 382 (1999); *J. Neurophysiol.* **82**, 398 (1999).
- N. Koshiya and J. C. Smith, *Nature* **400**, 360 (1999).
- H. Onimaru, A. Arata, I. Homma, *Jpn. J. Physiol.* **47**, 385 (1997).
- J. C. Reklung and J. L. Feldman, data not shown.
- E. Fletcher and J. Schaff, in *Abnormalities of Respiration During Sleep*, E. Fletcher, Ed. (Grune & Stratton, Orlando, FL, 1986), pp. 203–228; E. C. Fletcher, *Cardiologia* **42**, 469 (1997); S. Thalhofer and P. Dorow, *Respiration* **64**, 2 (1997).
- J. P. Orlowski, *Pediatrics* **78**, 233 (1986); V. Carpentier, H. Vaudry, E. Mallet, A. Laquerrière, P. Leroux, *Neuroscience* **86**, 159 (1998); A. Coquerel et al., *Neurochem. Int.* **20**, 97 (1992).
- Supported by a Ford Foundation Pre-Doctoral Fellowship for Minorities and a predoctoral fellowship from the Porter Physiology Development Program of the American Physiological Society to P.A.G., a Parker B. Francis Fellowship to J.C.R., and NIH grants HL40959 and HL37941. We thank N. Brecha, D. Buonomano, L. Kruger, and F. Schweizer for assistance with this manuscript.

12 July 1999; accepted 8 October 1999

Identification of a Conserved Receptor-Binding Site on the Fiber Proteins of CAR-Recognizing Adenoviridae

Peter W. Roelvink,* Gai Mi Lee, David A. Einfeld, Imre Kovcsdi, Thomas J. Wickham

The human adenovirus serotype 5 (Ad5) is used widely for applications in human gene therapy. Cellular attachment of Ad5 is mediated by binding of the carboxyl-terminal knob of its fiber coat protein to the Coxsackie adenovirus receptor (CAR) protein. However, Ad5 binding to CAR hampers the development of adenovirus vectors capable of specifically targeting (diseased) tissues or organs. Through sequence analysis and mutagenesis, a conserved receptor-binding region was identified on the side of three divergent CAR-binding knobs. The feasibility of simultaneous CAR ablation and redirection of an adenovirus to a new receptor is demonstrated.

The human adenovirus family has 49 viral serotypes that are associated with a wide range of pathologies and tissue tropisms (1, 2). The serotype Ad5 has been studied inten-

sively and is currently used as a vector in human gene therapy (3). Cellular attachment is mediated by binding of the COOH-terminal knob of the adenovirus fiber coat protein

LINKED CITATIONS

- Page 1 of 1 -



You have printed the following article:

Modulation of Respiratory Frequency by Peptidergic Input to Rhythmogenic Neurons in the PreBotzinger Complex

Paul A. Gray; Jens C. Reklung; Christopher M. Bocchiaro; Jack L. Feldman

Science, New Series, Vol. 286, No. 5444. (Nov. 19, 1999), pp. 1566-1568.

Stable URL:

<http://links.jstor.org/sici?sici=0036-8075%2819991119%293%3A286%3A5444%3C1566%3AMORFBP%3E2.0.CO%3B2-F>

This article references the following linked citations:

References and Notes

³ **Pre-Botzinger Complex: A Brainstem Region That May Generate Respiratory Rhythm in Mammals**

Jeffrey C. Smith; Howard H. Ellenberger; Klaus Ballanyi; Diethelm W. Richter; Jack L. Feldman
Science, New Series, Vol. 254, No. 5032. (Nov. 1, 1991), pp. 726-729.

Stable URL:

<http://links.jstor.org/sici?sici=0036-8075%2819911101%293%3A254%3A5032%3C726%3APCABRT%3E2.0.CO%3B2-M>

NOTE: *The reference numbering from the original has been maintained in this citation list.*