

Homologies in Leaf Form Inferred from *KNOX1* Gene Expression During Development

Geeta Bharathan,^{1*} Thomas E. Goliber,^{2*} Christopher Moore,^{2†} Sharon Kessler,² Thinh Pham,^{2‡} Neelima R. Sinha^{2§}

KNOTTED1-like homeobox (*KNOX1*) genes regulate development of the leaf from the shoot apical meristem (SAM) and may regulate leaf form. We examined *KNOX1* expression in SAMs of various vascular plants and found that *KNOX1* expression correlated with complex leaf primordia. However, complex primordia may mature into simple leaves. Therefore, not all simple leaves develop similarly, and final leaf morphology may not be an adequate predictor of homology.

In simple-leaved species (maize, rice, *Arabidopsis*, tobacco, snapdragon) *KNOX1* genes are expressed in the SAM and unexpanded axis and are down-regulated after leaf initiation, suggesting fundamental differences between the indeterminate shoot and determinate leaves (1–5). Overexpression of *KNOX1* in simple-leaved plants results in distorted leaves and ectopic shoots (6–8). By contrast, in tomato, a complex-leaved plant, *KNOX1* genes are up-regulated in leaf primordia and down-regulated in the mature leaf (9–11). Overexpression of *KNOX1* in tomato results in increased leaf complexity (9–11). To determine how general these patterns of expression are, we studied *KNOX1* expression in developing simple- and complex-leaved shoots in different species of a genus, *Lepidium*, and within a single species, *Neobeckia aquatica* (both in Brassicaceae, eurosid II).

All species showed down-regulation of *KNOX1* protein at sites of initiation of leaf primordia (P_0), but differed in whether or not *KNOX1* was expressed later in leaf development. The simple-leaved *Lepidium africanum* had high *KNOX1* protein expression in the SAM, but not later (Fig. 1, A and B), whereas in the complex-leaved *L. perfoliatum* and *L. hyssopifolium*, *KNOX1* expression appeared later in leaf primordia (Fig. 1, C to H). *L. oleraceum* produces simple leaves (Fig. 1I) and unexpectedly shows *KNOX1* expression in its leaf primordia (Fig. 1K) [see supplementary material (12)]. Scanning electron microscopy revealed that the simple leaves of *L. oleraceum* had primordia that, in early

development (primary morphogenesis), produced marginal outgrowths (Fig. 1J) typical of early complex leaf development (e.g., *L. hyssopifolium*: Fig. 1G) [see supplementary material (12)]. This early complex form of *L. oleraceum* was later subsumed by inner blade growth [by a process of secondary morphogenesis (13, 14)], and the early marginal outgrowths were apparent only as coarse teeth in the simple mature leaf (Fig. 1I). That *KNOX1* expression correlates with early leaf development but not necessarily with final leaf form implies that other regulatory changes influence leaf blade expansion.


The species of *Lepidium* studied here share an ancestor with complex leaves, as inferred from parsimony reconstructions of mature leaf form (Fig. 1L) on a phylogenetic hypothesis for the genus (15). Within this lineage simple leaves arose in a group that included *L. africanum* and *L. oleraceum*, and later, there was a reversal

back to complex leaves in *L. hyssopifolium* (Fig. 1L). We infer that the complex-leaved common ancestor had complex primordia with early *KNOX1* expression and that simple leaves evolved by either turning off *KNOX1* in the primordia (*L. africanum*) or modifying secondary morphogenesis (*L. oleraceum*) (16).

This correlation between *KNOX1* expression and primordium form was also observed within individuals of a single species, *N. aquatica* (Brassicaceae). This aquatic species has two kinds of leaves: simple or complex aerial leaves and submerged complex leaves. Emergent leaf form varies with light intensity: Simple leaves are produced under high light, and complex leaves under low light. *KNOX1* expression was absent in simple leaf primordia of emergent shoots under high light, but present in complex primordia made under low-light conditions (17). Thus, *KNOX1* expression can be modulated by light conditions, perhaps through hormonal changes that often accompany alterations in light quality and quantity (18).

Phylogenetic analyses of leaf evolution (Fig. 2) reveal that the ancestral angiosperm had simple leaves (19, 20), and that complex leaves repeatedly arose from these simple-leaved ancestors (on average 29 “gains”) and reverted (on average six “losses”) to the ancestral simple form [see supplemental material (12)]. This indicates that neither all simple nor all complex leaves are homologous [similar owing to common ancestry (21)]. Complex leaves are generally assumed to be nonhomologous (22), but simple leaves are generally assumed to be homologous and, therefore, developmentally similar. Our observations in *Lepidium* suggest that the latter assumption may not always be correct.

Fig. 1. Consistent correlation between leaf form and *KNOX* expression in Brassicaceae. (A and B) *L. africanum*; (C to E) *L. perfoliatum*; (F to H) *L. hyssopifolium*; (I to K) *L. oleraceum*. The final leaf form is shown in (A), (C), (F), and (I). Protein expression is present in (B), (D), (H), and (K) in the SAM (*); absent in P_0 and P_1 (O); and present in developing leaflets of complex leaf (Δ). The same symbols are used in Figs. 3 and 4. (E, inset) Whole-mount reverse transcription-polymerase chain reaction in situ hybridization shows expression of the *KNOX* gene *STM1* in developing leaflets. Scanning electron micrographs of *L. hyssopifolium* (G) and *L. oleraceum* (J) developing leaves. (L) Phylogenetic patterns of leaf evolution in *Lepidium* (Brassicaceae, eurosid I). Colors indicate ancestors reconstructed with simple (blue) or complex (red) leaves. Bars: (A, C, F, I) 1 cm, (G and J) 50 μ m, (B, D, H, K) 100 μ m. *KNOX* expression patterns: complex with no secondary simplification (asterisk), complex with secondary simplification (circled asterisk), or simple (filled yellow circle).



¹Department of Ecology and Evolution, State University of New York, Stony Brook, NY 11794–5245, USA.

²Section of Plant Biology, University of California–Davis, Davis, CA 95616, USA.

*These authors contributed equally to this work.

†Present address: Department of Genetics, Stanford University, Stanford, CA 94305, USA.

‡Present address: Department of Pathology, Genentech, South San Francisco, CA 94080, USA.

§To whom correspondence should be addressed. E-mail: nrsinha@ucdavis.edu

Does the correlation between *KNOX1* expression and primordium form hold for a broad range of taxa? We surveyed simple and complex leaves across vascular plants and found that (i) modification of complex primordia through secondary morphogenesis is common in simple leaves across eudicots; (ii) the presence of *KNOX1* expression is associated with complex

primordia and its absence with simple primordia; and (iii) *KNOX1* protein expression is down-regulated at the site of leaf initiation (P_0), except in ferns. Taxa were drawn from cycads, ferns, and various angiosperms that represent independent instances of origins of complex leaves or reversals to simple leaves.

KNOX1 protein expression reflective of simple leaves was seen in the simple primordia of *Amborella trichopoda* (Fig. 3, A to C), a putative basal extant angiosperm (23–25), and in the simple primordia of grasses (26, 27) [fig. S1 (12)]. Because primordia in basal angiosperms are simple (28), this expression pattern is inferred to represent the ancestral state in the

angiosperms. This inference is reinforced by our observation of the “simple” pattern in the simple-leaved gymnosperm, *Welwitschia* [fig. S1 (12)]. This is the pattern in most simple-leaved species studied to date (1–5) and in this study (*L. africanum*, *Neobeckia*, *Amborella*) and contrasts with the pattern seen in complex leaves of tomato [fig. S1 (12)]. However, as in *L. oleraceum*, development of simple leaves from complex primordia through secondary morphogenesis was also observed in various eudicot lineages [euasterid II: Apiales, *Pimpinella* (Fig. 3, D to F); euasterid I: Gentianales, *Coffea*; rosids: Vitales, *Vitis* sp. (fig. S2) (12)]. This molecular developmental dissimilarity may




Fig. 2. Evolution of leaf form. Summary of phylogenetic patterns inferred from parsimony reconstructions of ancestral states with data on leaf form in 557 genera of angiosperms (24, 39). The ancestral angiosperm (0), eudicot (1), rosids (2), and asterid (3) had simple leaves. Some groups at the tips (in red) are equally likely to have had ancestors with simple or complex leaves; within all other taxa in color there are multiple origins (“gains”) and reversals (“losses”) of complex leaves; taxa in black have only simple leaves. If reconstructions were done with terminals coded according to the state in the family (to include polymorphisms), then ancestors of taxa in green were reconstructed as having complex leaves. If polymorphic families were coded as having complex leaves, then the ancestral eudicot was equally likely to have had simple or complex leaves (1, closed red circle); if the ancestral eudicot is assumed to have had complex leaves, then a “loss” (4, open red circle) to simple leaves was followed by a new “gain” of complex leaves in the rosids (5, closed red circle). Under this scheme alone, it is possible that complex leaves of rosids (excluding Saxifragales) are homologous. *KNOX* expression patterns: as in Fig. 1.




Fig. 3. Comparison of mature leaf form, leaf primordia, and *KNOX1* immunolocalization pattern in angiosperms with simple leaves. “Simple” expression: (A to C) *A. trichopoda*. “Complex” expression: (D and E) *P. anisum* (Apiaceae). Scanning electron micrographs (B and E) show early development. *KNOX1* protein expression can be seen in shoot apices in (C) and (F). Symbols as in Fig. 1. Bars: (A and D) 1 cm, (B and E) 50 μm, (C and F) 100 μm.




Fig. 4. Vascular plants with complex pattern and complex leaves. (A and B) *D. carota* (Apiaceae). (C to E) *C. congestum* (Vitaceae). (F and G) *Anogramma chaephylla*. (H to J) *Zamia floridans*. Symbols as in Fig. 1. Bars: (A, C, F, H) 1 cm, (B, E, G, I), 100 μm, (D) 50 μm.

reflect the nonhomology of simple leaves in eudicots. KNOX1 expression of the "complex" pattern was seen in the complex primordia of different eudicot lineages [euasterid II: Apiaceae–*Daucus carota* (Fig. 4, A and B); rosids: Vitaceae–*Cissus* (Fig. 4, C to E)]. Sampling suggests that the molecular hypothesis regarding an association of KNOX1 activity with complexity of the leaf primordium is supported across most eudicots [fig. S3 (12)]. One exception is a group of legumes, including peas, which have complex leaves but no KNOX1 expression in leaf primordia (29, 30). We believe that, in this group, KNOX1 genes ceased to be part of the genetic cascade leading to the complex leaf form and that a different gene, PEAFL0 (29, 30), became part of the cascade (31). The unusual KNOX1 expression pattern in this group of legumes is striking, given our observation that the correlation of "complex" KNOX1 protein expression with primordium complexity was present in ferns and gymnosperms (Fig. 4, F to J), representing stages early in the evolution of vascular plants.

Regardless of final leaf form, KNOX1 expression is down-regulated at sites of leaf initiation (P_0) in flowering plants and gymnosperms [Figs. 1, 3, and 4; figs. S1 to S3 (12)]. This suggests a mechanism that denotes "determinacy" during initiation of the leaf. Unlike in seed plants, KNOX1 is not down-regulated in the P_0 of ferns (Fig. 4G) (32). This result is consistent with current understanding that leaves of ferns and seed plants evolved independently (33) and may have different developmental characteristics (34).

Our results suggest that at least two different modes of development have evolved to generate simple leaves (e.g., *L. africanum* with simple pattern and *Pimpinella anisum* with complex pattern). By contrast, the same, complex, pattern of KNOX1 expression characterizes independently evolved complex leaves (e.g., *Cissus congestum* and *Daucus carota*; *N. aquatica*, *L. perfoliatum*, and *L. hyssopifolium*). Complex leaves may thus be partially indeterminate. Several studies on vascular development in leaves suggest that leaf shape and venation patterns parallel each other. However, it is unclear whether one directs the other and, if so, which one. Analysis of venation patterns in developing leaves with secondary morphogenesis may provide some information on this aspect of leaf development. Our results are similar to those for Crustacea, a group of animals, in which Hox expression is correlated with the specialization of limbs into feeding appendages (35). As in that case, these results highlight the value of comparative studies in augmenting and/or refuting hypotheses that emerge from experimental studies, and in suggesting new hypotheses that may be tested experimentally.

References and Notes

1. E. Vollbrecht, B. Veit, N. Sinha, S. Hake, *Nature* **350**, 241 (1990).
2. C. Lincoln, J. Long, J. Yamaguchi, K. Serikawa, S. Hake, *Plant Cell* **6**, 1859 (1994).
3. A. Nishimura, M. Tamaoki, M. Matsuoka, *Plant Cell Physiol.* **39**, S60 (1998).
4. A. Nishimura, M. Tamaoki, Y. Sato, M. Matsuoka, *Plant J.* **18**, 337 (1999).
5. R. Waites, H. R. N. Selvadurai, I. R. Oliver, A. Hudson, *Cell* **93**, 779 (1998).
6. N. Sinha, R. E. Williams, S. Hake, *Genes Dev.* **7**, 787 (1993).
7. G. Chuck, C. Lincoln, S. Hake, *Plant Cell* **8**, 1277 (1996).
8. R. Schneeberger, M. Tsiantis, M. Freeling, J. A. Langdale, *Development* **125**, 2857 (1998).
9. D. Hareven, T. Gutfinger, A. Parnis, Y. Eshed, E. Lifschitz, *Cell* **84**, 735 (1996).
10. J.-J. Chen, B.-J. Janssen, A. Williams, N. Sinha, *Plant Cell* **9**, 1289 (1997).
11. B.-J. Janssen, L. Lund, N. Sinha, *Plant Physiol.* **117**, 771 (1998).
12. Supplementary figures and details of experimental procedures are available on Science Online at www.sciencemag.org/cgi/content/full/296/5574/1858/DC1.
13. W. Hagemann, S. Gleissberg, *Plant Syst. Evol.* **199**, 121 (1996).
14. N. G. Dengler, H. Tsukaya, *Int. J. Plant Sci.* **162**, 459 (2001).
15. J. L. Bowman, H. Bruggemann, J.-Y. Lee, K. Mummehoff, *Int. J. Plant Sci.* **160**, 917 (1999).
16. A recent phylogenetic hypothesis of relationships among many more *Lepidium* species on the basis of chloroplast sequences (36) suggests that these two shifts to simple leaves occurred independently. However, extensive hybridization may underlie this difference in phylogeny (37), and it is not possible to comment further on the independence or otherwise of this morphological shift.
17. N. R. Sinha et al., unpublished data.
18. J. Chory, J. Li, *Plant Cell Environ.* **20**, 801 (1997).
19. D. W. Taylor, L. J. Hickey, *Flowering Plant Origin, Evolution and Phylogeny* (Chapman & Hall, New York, 1996).
20. J. A. Doyle, P. K. Endress, *Int. J. Plant Sci.* **161**, S121 (2000).
21. Here we use the phylogenetic definition of homology. The present usage is prevalent in the field of evolution of development (38). Other uses of the term are not implied in this work.
22. S. Gleissberg, J. W. Kadereit, *Int. J. Plant Sci.* **160**, 787 (1999).
23. S. Mathews, M. Donoghue, *Science* **286**, 947 (1999).
24. Y.-L. Qiu et al., *Nature* **402**, 404, (1999).
25. D. E. Soltis et al., *Bot. J. Linn. Soc.* **133**, 381 (2000).
26. L. G. Smith, D. Jackson, S. Hake, *Dev. Genet.* **16**, 344 (1995).
27. Y. Sato et al., *Proc. Natl. Acad. Sci. U.S.A.* **93**, 8117 (1996).
28. W. Troll, *Vergleichende Morphologie der höheren Pflanzen Bd I. Vegetationsorgane* (Gebroeder Borntraeger, Berlin, 1939).
29. J. Hofer, L. Turner, R. Hellens, M. Ambrose, P. Mathews, *Curr. Biol.* **7**, 581, (1997).
30. C. W. Gourelay, J. M. I. Hofer, T. H. N. Ellis, *Plant Cell* **12**, 1279 (2000).
31. N. R. Sinha et al., unpublished data.
32. Similar results were observed in *Ceratopteris richardii* with a cloned KNOX1 gene (39).
33. P. Kenrick, P. R. Crane, *The Origin and Early Diversification of Land Plants: A Cladistic Study* (Smithsonian Institution Press, Washington, DC, 1997), pp. xiii–441.
34. Y. L. Ma, T. A. Steeves, *Ann. Bot.* **70**, 277 (1992).
35. M. Averof, N. H. Patel, *Nature* **388**, 682 (1997).
36. K. Mummehoff, H. Bruggemann, J. L. Bowman, *Am. J. Bot.* **88**, 2051 (2001).
37. K. Mummehoff, personal communication.
38. Abouheif et al., *Trends Genet.* **13**, 432 (1997).
39. J.-A. Banks, personal communication.
40. W. P. Maddison, D. R. Maddison, *MacClade: Interactive Analysis of Phylogeny and Character Evolution* (Sinauer Associates, Sunderland, MA, 1992).
40. We thank J. Harada, A. Doust, P. Stevens, and B. Grabowski for critical comments; T. Kellogg, C. Kuhlemeier, and members of the Sinha lab for helpful discussions; J. Jernstedt for discussions, help with dissections of the fern, cycad samples and, along with B. Hall, for the *Amborella* samples; T. Metcalf and E. Sandoval (Section of Plant Biology Conservatory) for plant materials; and S. Hake and D. Jackson for providing the antibodies to KNOTTED1. Supported by NSF IBN-9983063 and IBN-0092599 (N.R.S.) and SHARP (HHMI) undergraduate fellowships (C.M. and T.P.).

29 January 2002; accepted 15 April 2002

Amacrine-Signaled Loss of Intrinsic Axon Growth Ability by Retinal Ganglion Cells

Jeffrey L. Goldberg,* Matthew P. Klassen, Ying Hua, Ben A. Barres

The central nervous system (CNS) loses the ability to regenerate early during development, but it is not known why. The retina has long served as a simple model system for study of CNS regeneration. Here we show that amacrine cells signal neonatal rat retinal ganglion cells (RGCs) to undergo a profound and apparently irreversible loss of intrinsic axon growth ability. Concurrently, retinal maturation triggers RGCs to greatly increase their dendritic growth ability. These results suggest that adult CNS neurons fail to regenerate not only because of CNS glial inhibition but also because of a loss of intrinsic axon growth ability.

Neurons in the CNS lose the ability to regenerate their axons early in development, but it is not known why. A currently prevailing view is that a strongly inhibitory glial environment causes regenerative failure in the adult CNS (1, 2), as CNS glial cells, both astrocytes and oligodendrocytes, inhibit re-

generating axons after injury (3–6). A crucial question is whether overcoming these inhibitory cues will be sufficient to promote rapid regeneration or whether adult CNS neurons have undergone a developmental loss of intrinsic regenerative ability (7–11). For example, CNS neurons in slices are less able to