REPORTS

- 20. T. R. McGetchin, M. Settle, J. W. Head, Earth Planet. Sci. Lett. 20, 226 (1973).
- 21. D. Stoffler, D. E Gault, J. Wedekind, G. Polkowski, J. Geophys. Res. 80, 29 (1975).
- 22. P. E. Olsen, D. V. Kent, R. Raeside "International workshop for a climatic, biotic, and tectonic, pole-topole coring transect of Triassic-Iurassic Pangea" (International Continental Scientific Drilling Program, Potsdam, 1999), vol. 1; www.ldeo.columbia.edu/ ~polsen/nbcp/icdp.abst.html.
- 23. M. W. Hounslow, G. Warrington, P. E. Posen, poster presented at the International Geological Correlation Programme (IGCP) 458 Southwest England Field Workshop: Jurassic/Triassic Boundary Meeting, Taunton, UK, October 2001; http://geography.lancs.ac.uk/ cemp/publications/01houns-sta_abs.htm
- 24. D. M. Bice, C. R. Newton, S. McCauley, P. W. Reiners, C. A. McRoberts, Science 225, 443 (1992).
- 25. We thank I. Whitehead for his guidance on measuring planar deformation features in quartz using a u-stage.

Heart Regeneration in Zebrafish

Kenneth D. Poss,* Lindsay G. Wilson, Mark T. Keating*

Cardiac injury in mammals and amphibians typically leads to scarring, with minimal regeneration of heart muscle. Here, we demonstrate histologically that zebrafish fully regenerate hearts within 2 months of 20% ventricular resection. Regeneration occurs through robust proliferation of cardiomyocytes localized at the leading epicardial edge of the new myocardium. The hearts of zebrafish with mutations in the Mps1 mitotic checkpoint kinase, a critical cell cycle regulator, failed to regenerate and formed scars. Thus, injury-induced cardiomyocyte proliferation in zebrafish can overcome scar formation, allowing cardiac muscle regeneration. These findings indicate that zebrafish will be useful for genetically dissecting the molecular mechanisms of cardiac regeneration.

Injured human hearts do not regenerate. Instead, damaged myocardium is replaced by fibrotic scar tissue. Cardiomyocytes, the major structural cells of the heart, may undergo hypertrophy in the wound area to increase muscular mass. Although recent findings suggest

Supporting Online Material

www.sciencemag.org/cgi/content/full/1076249/DC1 Figs. S1 and S2 Tables S1 and S2 References

17 July 2002; accepted 5 November 2002 Published online 14 November 2002; 10.1126/science.1076249 Include this information when citing this paper.

that cardiomyocytes within the diseased human heart can proliferate (1), most evidence to date indicates that myocyte proliferation is not a significant component of the mammalian response to cardiac injury (2).

Teleost fish, including zebrafish, can regenerate spinal cord, retina, and fins (3, 4). To determine whether zebrafish can also regenerate heart muscle, we surgically removed \sim 20% of the ventricular myocardium from 1to 2-year-old adults (Fig. 1, A and B) and

Department of Cell Biology, Department of Cardiology, Howard Hughes Medical Institute, Harvard Medical School, Children's Hospital, 320 Longwood Avenue, Boston, MA 02115, USA.

*To whom correspondence should be addressed. E-mail: kposs@enders.tch.harvard.edu(K.D.P.); mkeating@enders. tch.harvard.edu (M.T.K.)



sealed with a large amount of mature fibrin. (F) 14 dpa. The fibrin has diminished, and the heart muscle has reconstituted. (G) 30 dpa. A new cardiac wall has been created, and only a small amount of internal fibrin remains (arrowhead). (H) 60 dpa. This ventricle shows no sign of injury. (I) Quantification of healing at 0, 30, and 60 dpa. Values represent the size of the largest ventricular section (mean \pm SEM; *P < 0.05); parentheses indicate the number of hearts examined (5). Scale bars, 100 µm.



ximate

examined hearts histologically at eight time points after surgery (5). After several seconds of profuse bleeding from the ventricular lumen, a large clot of erythrocytes formed in the wound (Fig. 1, C and D). Beginning at 2 to 4 days postamputation (dpa), these erythrocytes were replaced by fibrin, which reached maximum levels at 7 to 9 dpa (Fig. 1E). For the first few days after ventricular resection, zebrafish appeared less active and less coordinated while swimming than shamoperated animals. After 1 week of recovery, however, they were indistinguishable from controls.

From 9 to 30 dpa, cardiac myofibers surrounded, penetrated, and eventually replaced the clot (Fig. 1, F and G). By 30 dpa, a contiguous wall of muscle had formed (Fig. 1G); by 60 dpa, the fibrin clot had completely

Fig. 2. Cardiomyocyte proliferation accompanies zebrafish heart regeneration. (A) Confocal image of a heart section of an unamputated fish labeled for 7 days with BrdU, stained for myosin heavy chain to identify cardiomyocytes (red), and stained with BrdU to detect cycling cells (green) and with 4',6'-diamidino-2-phenylindole (DAPI) to detect nuclei (blue). A low percentage of compact myocytes incorporate BrdU over this period (arrowhead) (5). (B) 7 dpa (0- to 7-dpa BrdU labeling). BrdU incorporation occurs in trabecular myocytes at the amputation plane (arrowheads). Most hearts also showed labeling in compact myocytes adjacent to the wound area at this stage (not shown). (C) 14 dpa (7- to 14-dpa BrdU labeling). Many cardiomyocytes incorporate BrdU during this period, largely in compact muscle adjacent to the wound. (D) 30 dpa (23- to 30-dpa BrdU labeling). Myocyte BrdU incorporation continues as the compact muscular laver expands. Labeldisappeared, and the size and shape of most zebrafish ventricles appeared grossly normal by histology (Fig. 1H) (5). The contractile properties of beating hearts in situ at 60 dpa also appeared grossly normal through direct visual inspection (δ).

To quantify the replacement of zebrafish heart muscle, we measured the surface area of longitudinal ventricular sections of injured hearts by digital imaging and computer quantification. Amputation removed ~19% of the ventricular section surface area (Fig. 1I). Measurements taken 30 dpa revealed partial recovery of ventricular section surface area, and, by 60 dpa, the surface area was completely recovered (Fig. 1I) (5).

The zebrafish ventricle is composed of a thin, external layer of compact myocytes penetrated by blood vessels and by internal myo-



ing is usually limited to the most epicardial myocytes of the apex (arrowheads). (E) Cardiomyocyte mitoses observed at 14 dpa in the wound area, with the use of an antibody to phosphorylated histone-3 (H3P; arrowheads indicate positive cells). Insets show a cardiomyocyte with sister chromatid segregation (left) and cardiomyocytes that have recently completed karyokinesis (right). Sections are stained for myosin heavy chain (red) and H3P (green) and stained with DAPI (blue). (F) 30 dpa (7- to 14-dpa BrdU labeling). Arrows indicate the area of incorporation. There is an increase in labeled cardiomyocytes in 30-dpa pulse-chase hearts. Several unlabeled layers of new myocytes are external to labeled myocytes, presumably resulting from cell division after the labeling period. Scale bars, 25 μ m; 5 μ m in (E) insets.

cytes organized into elaborate trabeculae (fig. S1). An antibody to β catenin, which is localized to adherens junctions (7, 8), clearly demarcated these two types of myocardium. This method revealed that the initial wound area through 7 dpa consisted mainly of trabecular myocytes exposed to the clot. By 14 dpa, most of the wound was enveloped by a wall of compact myocytes, leaving only a small portion of medial trabecular myocardium exposed to the clot. By 30 dpa, and even more prominently at 60 dpa, a large, contiguous wall of compact muscle 10 to over 30 myocytes wide was created at the apex, replacing the resected compact and trabecular myocardium (fig. S1). These observations indicate that a new layer of compact myocytes was reestablished and then expanded after amputation.

The restoration of cardiac muscle could result from myocyte hypertrophy or hyperplasia. To assess zebrafish cardiomyocyte proliferation, we assayed cell cycle entry by measuring the nuclear incorporation of bromodeoxyuridine (BrdU), a marker of DNA synthesis. In uninjured hearts, about 3% of compact cardiomyocyte nuclei incorporated BrdU after a 7-day labeling period, with no trabecular cardiomyocyte incorporation (Fig. 2A and fig. S2). In contrast, in injured hearts \sim 17% of compact and trabecular myocytes near the amputation plane incorporated BrdU at 7 dpa (Fig. 2B). BrdU incorporation peaked at \sim 32% at 14 dpa, with most cycling myocytes localized to compact muscle at the lateral edges of the wound (Fig. 2C). At 30 dpa, the BrdU incorporation index in compact muscle was still 20% (Fig. 2D) but had considerably decreased to 7% (fig. S2) by 60 dpa. Cardiac myocytes also underwent mitosis, as defined by expression of phosphorylated histone-3, a marker of condensed chromatin. Although mitoses were rarely seen in the uninjured heart, we observed 3 to 10 cardiomyocyte mitoses per wound area at 14 dpa (Fig. 2E).

We next gave a pulse of BrdU to zebrafish that were from 7 to 14 dpa and assessed cardiomyocyte labeling at 14, 30, and 60 dpa. After the pulse, BrdU-positive cells continued to increase in number and were often found adjacent to one another in the regenerated myocardium, consistent with cardiomyocyte cell division (Fig. 2F). Nuclei in these myocytes were frequently punctate-labeled, suggesting BrdU dilution via karyokinesis (fig. S3).

Pulse-chase experiments also revealed that the leading edge of proliferation during regeneration was the new layer of outermost (epicardial) myocytes. Zebrafish labeled with BrdU at 30 and 60 dpa demonstrated proliferation in epicardial cardiomyocytes of the new compact muscle (Fig. 2D). By contrast, zebrafish labeled from 7 to 14 dpa with BrdU and given a 16- or 46-day chase showed substantial BrdU incorporation in the inner layers of the new compact myocardium, with extremely limited labeling in epicardial myocytes (Fig. 2F and fig. S3). These data indicate that, after dispersed cardiomyocyte proliferation in the wound area to restore the ventricular wall, zebrafish cardiac regeneration occurs via a gradient of proliferation that is greatest in epicardial myocytes. Regenerated myocytes are displaced inward as epicardial myocytes continue to proliferate and advance.

Fibrosis is the dominant histological reaction to injury in mammalian and amphibian hearts. This response typically follows fibrin deposition after cardiac injury (9). By contrast, zebrafish ventricles displayed only small deposits of collagen extending into the fibrin clot at 14 dpa (Fig. 3A). The extent of collagen deposition at 30 and 60 dpa varied from complete absence to small deposits (n =13 hearts) (Fig. 3B).

Why do zebrafish respond to cardiac injury with regeneration, whereas fibrosis predominates in other vertebrates? We propose a model in which scarring complements regeneration, and it is the vigor of myocyte proliferation within a given species that determines the predominant response. This model predicts that inhibition of regeneration would lead to scarring. To test this model, we ana-

Fig. 3. Cardiomyocyte proliferation is required for scarless regeneration. 14-dpa (A) and 60dpa (B) ventricles stained with acid fuchsin-orange G (AFOG; fibrin, orange/red; collagen, blue), which is highly sensitive for collagen (arrowheads). Regenerated ventricles contain minimal collagen. (C and D) 17-dpa ventricles double-stained for myosin heavy chain and with aniline blue (muscle, brown; fibrin and collagen, grayish blue). Wild-type ventricles display new compact muscular wall formation (C), whereas mps1 mutant ventricles demonstrate no evidence of new muscle (D) (14). (E and F) 26-dpa hearts stained with AFOG. The wild-type cardiac injury response includes minor fibrin retention and collagen deposition (E), whereas extensive fibrosis (arrowheads) is observed in cardiac wounds of mps1 mutants (F). Scale bars, 100 µm.

lyzed the cardiac injury response in mps1 mutant zebrafish. These animals harbor a temperature-sensitive mutation in Mps1, a mitotic checkpoint kinase that is up-regulated in many proliferative cell types (10-13). At the restrictive temperature, this mutation potently blocks fin regeneration through failed cell proliferation (13). In situ hybridization of 14-dpa zebrafish hearts showed induction of mps1 mRNA in a small number of myocytes near the wound, consistent with localization and function of Mps1 in regenerating cardiomyocytes (fig. S4). The mps1 mutants formed normal fibrin clots by 8 to 9 dpa (6). In wild-type fish, cardiac myofibers invariably penetrated the clot and constructed a bridge of new muscle around the wound by 17 to 26 dpa (Fig. 3C) (n = 12). By contrast, the ventricular wall was not restored in mps1 mutants (Fig. 3D) (n = 13); instead, the injured hearts retained fibrin deposits and developed large connective-tissue scars (Fig. 3, E and F) (n = 7) (14).

We conclude that, in response to mechanical injury, zebrafish hearts can regenerate without scarring. A new ventricular wall of compact myocardium is created in the injured heart by cardiomyocyte proliferation, a process that requires the cell-cycle regulator Mps1. Although cardiomyocyte proliferation in injured adult amphibian hearts has been documented (15-19), we suggest that ze-



brafish represent an improved model system for understanding heart regeneration. Zebrafish restore the ventricular myocardial wall, whereas published findings indicate little or no cardiac muscle regeneration after amphibian heart resection. Also, a wide array of molecular and genetic tools, including mutagenesis screens and complete genome sequence information, are available to those working with zebrafish. Thus, dissection of zebrafish cardiac muscle regeneration may illuminate factors that can stimulate a regenerative response in the mammalian heart.

References and Notes

- 1. A. P. Beltrami *et al.*, *N. Engl. J. Med.* **344**, 1750 (2001).
- 2. K. B. Pasumarthi, L. J. Field, Circ. Res. 90, 1044 (2002).
- S. L. Johnson, J. A. Weston, *Genetics* 141, 1583 (1995).
- T. Becker, M. F. Wullimann, C. G. Becker, R. R. Bernhardt, M. Schachner, J. Comp. Neurol. 377, 577 (1997).
- Materials and methods are available as supplemental material on Science Online.
- K. D. Poss, L. G. Wilson, M. T. Keating, unpublished observations.
- 7. T. Kurth, H. Schwarz, S. Schneider, P. Hausen, *Cell Tissue Res.* 286, 1 (1996).
- K. K. Linask, K. A. Knudsen, Y. H. Gui, Dev. Biol. 185, 148 (1997).
- 9. S. J. Schnitt et al., Circ. Res. 72, 914 (1993).
- E. Weiss, M. Winey, J. Cell Biol. 132, 111 (1996).
 A. Abrieu et al., Cell 106, 83 (2001).
- A. Abrieu et al., Cell 106, 85 (2001).
 D. Hogg et al., Oncogene 9, 89 (1994).
- K. D. Poss, A. Nechiporuk, A. M. Hillam, S. L. Johnson,
- M. T. Keating, Development 129, 5141 (2002). 14. The mos1 mutants and wild-type controls were
- maintained at 25°C before and for 24 hours after surgery and then shifted to 32° to 33°C-recirculating aquarium water for 7 to 25 days. Pulse-chase studies with BrdU showed no evidence of myocyte proliferation in *mps1* mutants during this period. Animals of the WIK or (WIK × *AB) strains, closest in genetic background to the *mps1* mutant strain, were used as wild-type controls. The WIK strain had more difficulty surviving surgery than other strains. So, in these experiments, we removed only 10 to 15% of the ventricle. Regeneration appears to occur less effectively at 32° to 33°C than at 25°C in wild-type fish, with less myocyte proliferation (6).
- J. O. Oberpriller, J. C. Oberpriller, J. Exp. Zool. 187, 249 (1974).
- 16. D. Bader, J. O. Oberpriller, J. Morphol. 155, 349 (1978).
- P. P. Rumyantsev, Z. Zellforsch Mikrosk. Anat. 139, 431 (1973).
- A. W. Neff, A. E. Dent, J. B. Armstrong, Int. J. Dev. Biol. 40, 719 (1996).
- 19. I. L. Flink, Anat. Embryol. 205, 235 (2002).
- 20. We thank A. Hillam for help with histology; A. Nechiporuk and S. Johnson for helping to generate and analyze mps1 mutants; H. Stern and R. Bronson for help with histopathology; D. Clapham for confocal imaging equipment; A. Nechiporuk, S. Odelberg, A. Limbourg, E. Lien, and L. Zon for critique of the manuscript; A. Sanchez and S. Kim for animal care; and Keating laboratory members for helpful discussions. Supported by the Helen Hay Whitney Foundation (K.D.P.) and grants from the National Heart, Lung, and Blood Institute. All animal procedures were performed in accordance with Children's Hospital guidelines.

Supporting Online Material

www.sciencemag.org/cgi/content/full/298/5601/2188/

Materials and Methods Figs. S1 to S4

28 August 2002; accepted 23 October 2002